

Extraction of Meson PDFs from Drell-Yan and J/ψ Production Data in the Statistical Model

Jen-Chieh Peng

University of Illinois at Urbana-Champaign

The 12th Circum-Pan-Pacific Symposium on High Energy
Spin Physics, Nov. 8-13, 2024

Hefei, China

In collaboration with Claude Bourrely (Marseille),
Franco Buccella (Rome), and Wen-Chen Chang (Taipei)

Partonic structures of pion and kaon

Why is it interesting?

- Lightest $q\bar{q}$ bound states, and Goldstone bosons
- A simpler hadronic system than the nucleon
- Provide information on mass decomposition of pion and kaon
- Spin-0 π and K contrasting spin-1/2 nucleon
- Compared to nucleons, very little is known experimentally for the partonic structures of mesons

Partonic structures of pion and kaon

Spin-0 for π and K implies:

- No helicity distributions ($\Delta q(x) = 0$, $\Delta G(x) = 0$)
- No TMDs such as Transversity, Sivers, Pretzelosity distributions (Boer-Mulders functions for π and K do exist)

Number of unpolarized partonic distributions is reduced from symmetry consideration (charge-conjugation (C) and SU(2) flavor (I) symmetries)

$$\bullet u_{\pi^+}^V(x) \stackrel{C}{=} \bar{u}_{\pi^-}^V(x) \stackrel{I}{=} \bar{d}_{\pi^+}^V(x) \stackrel{C}{=} d_{\pi^-}^V(x) \equiv V_{\pi}(x)$$

$$\bullet \bar{u}_{\pi^+}(x) \stackrel{C}{=} u_{\pi^-}(x) \stackrel{I}{=} d_{\pi^+}(x) \stackrel{C}{=} \bar{d}_{\pi^-}(x) \equiv S_{\pi}(x)$$

For kaons, more PDFs are needed (breaking of SU(3) flavor symmetry)

$$\bullet u_{K^+}^V(x) \neq \bar{s}_{K^+}^V(x) \text{ (analogous to } u_p^V(x) \neq d_p^V(x))$$

$$\bullet \bar{u}_{K^+}(x) \neq \bar{d}_{K^+}(x) \text{ (analogous to } \bar{u}_p(x) \neq \bar{d}_p(x))$$

Many interesting questions can be raised on the comparison between pion and kaon parton distributions


Meson partonic content from the Drell-Yan Process

MASSIVE LEPTON-PAIR PRODUCTION IN HADRON-HADRON COLLISIONS AT HIGH ENERGIES*

Sidney D. Drell and Tung-Mow Yan

Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305

(Received 25 May 1970)



$$p + p \rightarrow (\mu^+ \mu^-) + \dots \quad (1)$$

Our remarks apply equally to any colliding pair such as (pp) , $(\bar{p}p)$, (πp) , (γp) and to final leptons $(\mu^+ \mu^-)$, $(e\bar{e})$, $(\mu\nu)$, and $(e\nu)$.

(4) The full range of processes of the type (1) with incident p , \bar{p} , π , K , γ , etc., affords the interesting possibility of comparing their parton and antiparton structures.

Drell-Yan experiments with π^- beam

Experiments at CERN and Fermilab

Exp	P (GeV)	targets	Number of D-Y events
WA11	175	Be	500 (semi-exclusive)
WA39	40	W (H ₂)	3839 (all beam, M > 2 GeV)
NA3	150, 200, 280	Pt (H ₂)	21600, 4970, 20000 (535, 121, 741)
NA10	140, 194, 286	W (D ₂)	~84400, ~150000, ~45900 (3200, -, 7800)
E331/E44 4	225	C, Cu, W	500
E326	225	W	
E615	80, 252	W	4060, ~50000

- Relatively pure π^- beam; J/ Ψ production also measured
- Relatively large cross section due to $\bar{u}d$ contents in π^-

For a very long time, only four pion parton distribution functions were available

- First: OW-P (PRD 30, 943 (1984))

- Second: ABFKW-P (PL 233, 517 (1989))

- Third: GRV-P (Z. Phys. C35, 651 (1992))

- Fourth: SMRS (PR D45, 2349 (1992))


- Need new global fits to all existing data
- Need new experimental data with pion and kaon beams

First Monte Carlo global QCD analysis of pion parton distributions

P. C. Barry,¹ N. Sato,² W. Melnitchouk,³ and Chueng-Ryong Ji¹

JAM Collaboration

PRL 121, 152001 (2018);
PRL 127, 232001 (2021)



- Drell-Yan data from NA10 and E615
- **Leading-neutron tagged DIS from HERA** provides information on the pion PDFs at small x
- The Q^2 evolution allows extraction of gluon distribution
- Uncertainties of the pion PDFs are determined

**Parton distribution functions of the charged pion
within the xFitter framework**

Ivan Novikov^{1,2,*}, Hamed Abdolmaleki³, Daniel Britzger⁴, Amanda Cooper-Sarkar⁵, Francesco Giuli⁶,
Alexander Glazov^{2,†}, Aleksander Kusina⁷, Agnieszka Luszczak⁸, Fred Olness⁹, Pavel Starovoitov¹⁰,
Mark Sutton¹¹ and Oleksandr Zenaiev¹²

(xFitter Developers' team)

- Drell-Yan data from NA10 and E615
- Direct photon production data from WA70
- Uncertainties of the pion PDFs are determined
- Valence distribution is well determined, but not the sea and gluon distributions

A New Extraction of Pion Parton Distributions in the Statistical Model

Claude Bourrely^a, Franco Buccella^b, Jen-Chieh Peng^c

Physics Letters B 813 (2021) 136021

$$xU(x) = xD(x) = \frac{A_U X_U x^{b_U}}{\exp[(x - X_U)/\bar{x}] + 1} + \frac{\tilde{A}_U x^{\tilde{b}_U}}{\exp(x/\bar{x}) + 1}. \quad (7)$$


$$x\bar{U}(x) = x\bar{D}(x) = \frac{A_U (X_U)^{-1} x^{b_U}}{\exp[(x + X_U)/\bar{x}] + 1} + \frac{\tilde{A}_U x^{\tilde{b}_U}}{\exp(x/\bar{x}) + 1}. \quad (8)$$

$$xS(x) = x\bar{S}(x) = \frac{\tilde{A}_U x^{\tilde{b}_U}}{2[\exp(x/\bar{x}) + 1]}. \quad (9)$$


$$xG(x) = \frac{A_G x^{b_G}}{\exp(x/\bar{x}) - 1}. \quad (10)$$

- The statistical model describes proton's PDF very well
- The antiquark's flavor structure is related to quark's flavor structure
- The antiquark's spin structure is related to quark's spin structure
- It is not clear if the statistical model also works for meson's PDFs


E615 $\pi^- W \rightarrow \mu^- \mu^+ X$ 252 GeV



NA10 $\pi^- W \rightarrow \mu^- \mu^+ X$ 194 GeV



NA10 $\pi^- W \rightarrow \mu^- \mu^+ X$ 286 GeV



With only a few parameters for the pion PDFs, the Drell-Yan data are well described by the statistical model

Comparison between proton and pion PDFs in the statistical model

$$xQ^\pm(x) = \frac{A_Q X_Q^\pm x^{b_Q}}{\exp[(x - X_Q^\pm)/\bar{x}] + 1},$$

$$A_U = 0.776 \pm 0.15$$

$$b_U = 0.500 \pm 0.02$$

$$X_U = 0.756 \pm 0.01$$

$$\bar{x} = 0.1063 \pm 0.004$$

$$\tilde{A}_U = 2.089 \pm 0.21$$





$$\tilde{b}_U = 0.4577 \pm 0.009$$

$$A_G = 31.17 \pm 1.7$$

$$b_G = 1 + \tilde{b}_U.$$

- The temperature, $\bar{x} = 0.106$, found for pion is very close to that obtained for proton, $\bar{x} = 0.090$, suggesting a common feature for the statistical model description of baryons and mesons
- The chemical potential of the valence quark for pion, $X_U = 0.756$, is significantly larger than for proton, $X_U = 0.39$

Valence and gluon distributions for various pion PDFs




- Quite good agreements for valence quark PDFs
- Much larger variations for the gluon PDFs


Constraining gluon distribution of pion with pion-induced J/Ψ production

- The Drell-Yan data are not sensitive to the gluon distributions in pion
- The J/Ψ production data are sensitive to the gluon PDF in pion, which is poorly known and is of much theoretical interest

J/Ψ (q - q bar annihilation)



J/Ψ (gluon-gluon fusion)



Pion PDFs using DY and J/Ψ data in the statistical model

PHYSICAL REVIEW D **105**, 076018 (2022)

Pion partonic distributions in a statistical model from pion-induced Drell-Yan and J/Ψ production data

Claude Bourrely¹, Wen-Chen Chang², and Jen-Chieh Peng³

¹*Aix Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France*

²*Institute of Physics, Academia Sinica, Taipei 11529, Taiwan*

³*Department of Physics, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA*



(Received 23 February 2022; accepted 6 April 2022; published 26 April 2022)

We present a new analysis to extract pion parton distribution functions (PDFs) within the framework of the statistical model. Starting from the statistical model first developed for the spin-1/2 nucleon, we extend this model to describe the spin-0 pion. Based on a combined fit to both the pion-induced Drell-Yan data and the pion-induced J/Ψ production data, a new set of pion PDFs has been obtained. The inclusion of the J/Ψ production data in the combined fit has provided additional constraints for better determining the gluon distribution in the pion. We also compare the pion PDFs obtained in the statistical model with other existing pion PDFs.

Pion PDFs using DY and J/Ψ data

Phys.Rev.D 105 (2022) 076018 ; arXiv: 2202.12547



$$xU(x) = xD(x) = \frac{A_U X_U x^{b_U}}{\exp[(x - X_U)/\bar{x}] + 1} + \frac{\tilde{A}_U x^{\tilde{b}_U}}{\exp(x/\bar{x}) + 1}$$

$$x\bar{U}(x) = x\bar{D}(x) = \frac{A_U (X_U)^{-1} x^{b_U}}{\exp[(x + X_U)/\bar{x}] + 1} + \frac{\tilde{A}_U x^{\tilde{b}_U}}{\exp(x/\bar{x}) + 1}$$

$$xS(x) = x\bar{S}(x) = \frac{\tilde{A}_U x^{\tilde{b}_U}}{2[\exp(x/\bar{x}) + 1]}$$

$$xG(x) = \frac{A_G x^{b_G}}{\exp(x/\bar{x}) - 1}, \quad b_G = 1 + \tilde{b}_U$$

J / Ψ WA39 and NA3 J/ψ (π⁻ H₂)




Inclusion of the J/Ψ data gives larger G(x) at x>0.1

NRQCD for J/Ψ Production

What do we know about the kaon PDF (very little!)

$\sigma(K^- + Pt) / \sigma(\pi^- + Pt)$ Drell-Yan ratios



From NA3; 150 GeV, Pt target

$$R = \frac{\sigma_{DY}(K^- + D)}{\sigma_{DY}(\pi^- + D)}$$



$$\simeq \frac{4V_K^u(x_1)V_N(x_2) + 4V_K^u(x_1)S_N(x_2) + V_K^s(x_1)s_p(x_2) + 5S_K(x_1)V_N(x_2)}{4V_\pi(x_1)V_N(x_2) + 5S_\pi(x_1)V_N(x_2) + 5V_\pi(x_1)S_N(x_2)} \simeq \frac{V_K^u(x_1)}{V_\pi(x_1)}$$

$R \simeq (1-x)^{0.18 \pm 0.07} \Rightarrow$ softer u -valence in kaon than in pion

$(K^- + Pt) / (\pi^- + Pt)$ ratios for J/Ψ production

From NA3; 150 GeV, Pt target

Ratios for D-Y



Similar behavior at large x_F for D-Y and J/Ψ production?

Extraction of kaon partonic distribution functions from Drell-Yan and J/ψ production data

Claude Bourrely ^{a, ID, *}, Franco Buccella ^b, Wen-Chen Chang ^c, Jen-Chieh Peng ^d

Pion PDFs

Phys. Lett. B 848 (2024) 138395

Kaon PDFs

$$xU_{\pi}(x) = \frac{A_U X_U x^{b_U}}{\exp[(x - X_U)/\bar{x}] + 1} + \frac{\tilde{A}_U x^{\tilde{b}_U}}{\exp(x/\bar{x}) + 1} ;$$

$$x\bar{U}_{\pi}(x) = \frac{A_U (X_U)^{-1} x^{b_U}}{\exp[(x + X_U)/\bar{x}] + 1} + \frac{\tilde{A}_U x^{\tilde{b}_U}}{\exp(x/\bar{x}) + 1} ;$$

$$xS_{\pi}(x) = \frac{\tilde{A}_U x^{\tilde{b}_U}}{2[\exp(x/\bar{x}) + 1]} ;$$

$$xG_{\pi}(x) = \frac{A_G x^{b_G}}{\exp(x/\bar{x}) - 1}, \quad b_G = 1 + \tilde{b}_U .$$

$$xU_K(x) = \frac{A_{UK} X_{UK} x^{b_{UK}}}{\exp[(x - X_{UK})/\bar{x}] + 1} + \frac{\tilde{A}_{UK} x^{\tilde{b}_{UK}}}{\exp(x/\bar{x}) + 1} ;$$


$$x\bar{U}_K(x) = \frac{A_{UK} (X_{UK})^{-1} x^{b_{UK}}}{\exp[(x + X_{UK})/\bar{x}] + 1} + \frac{\tilde{A}_{UK} x^{\tilde{b}_{UK}}}{\exp(x/\bar{x}) + 1} ;$$

$$xS_K(x) = \frac{A_{SK} X_{SK} x^{b_{SK}}}{\exp[(x - X_{SK})/\bar{x}] + 1} + \frac{\tilde{A}_{UK} x^{\tilde{b}_{UK}}}{2[\exp(x/\bar{x}) + 1]} ;$$

$$x\bar{S}_K(x) = \frac{A_{SK} (X_{SK})^{-1} x^{b_{SK}}}{\exp[(x + X_{SK})/\bar{x}] + 1} + \frac{\tilde{A}_{UK} x^{\tilde{b}_{UK}}}{2[\exp(x/\bar{x}) + 1]} ;$$


$$xD_K(x) = x\bar{D}_K(x) = \frac{\tilde{A}_{UK} x^{\tilde{b}_{UK}}}{(\exp(x/\bar{x}) + 1)} ;$$

$$xG_K(x) = \frac{A_{GK} x^{b_{GK}}}{\exp(x/\bar{x}) - 1}, \quad b_{GK} = 1 + \tilde{b}_{UK} .$$




The K^- / π^- D-Y data
can be well described

Extraction of kaon partonic distribution functions from Drell-Yan and J/ψ production data

Claude Bourrely^{a, *}, Franco Buccella^b, Wen-Chen Chang^c, Jen-Chieh Peng^d


Phys. Lett. B 848 (2024) 138395



The K^-/π^- and K^+/π^+ J/ψ data can also be well described by the statistical model (red curves)


The dashed curves use the recent PDFs obtained in the "Maximum Entropy" approach

Extraction of kaon partonic distribution functions from Drell-Yan and J/ψ production data


Claude Bourrely ^{a, , *}, Franco Buccella ^b, Wen-Chen Chang ^c, Jen-Chieh Peng ^d

Phys. Lett. B 848 (2024) 138395

Comparison between the pion and kaon valence distributions



Comparison between the pion and kaon gluon distributions



Momentum fractions of valence quarks, sea quarks, and gluons for π^- and K^- at the scale $Q^2 = 10 \text{ GeV}^2$ obtained in the statistical model.


	u Valence	d Valence	s Valence	all Sea	Gluon
π^-	0.242 ± 0.004	0.242 ± 0.004	–	0.188 ± 0.004	0.326 ± 0.015
K^-	0.220 ± 0.002	–	0.276 ± 0.001	0.162 ± 0.006	0.331 ± 0.018

$$S_K > U_\pi > U_K; \quad G_K \simeq G_\pi$$

More Drell-Yan and J/ψ data are coming

New pion-induced Drell-Yan data from COMPASS

Drell-Yan cross section on W and comparison to E615



$$\sqrt{\tau} = M/\sqrt{s}$$


- **New results since 30 years**
- Similar kinematic coverage as E615
- Better statistics, similar total systematics except for the low mass region



More Drell-Yan and J/ψ data are coming


New pion-induced Drell-Yan data from COMPASS

3 dimensional Drell-Yan cross section on $\text{NH}_3\text{-He}$



- First high statistics measurement with light material
- Red line/shaded area: statistical / total (stat. and syst.) uncertainties
- Dominated by statistical uncertainty

AMBER (Phase-I was approved)



Program	Physics Goals	Beam Energy [GeV]	Beam Intensity [s^{-1}]	Trigger Rate [kHz]	Beam Type	Target	Earliest start time, duration
muon-proton elastic scattering	Precision proton-radius measurement	100	$4 \cdot 10^6$	100	μ^\pm	high-pressure H2	2022 1 year
Hard exclusive reactions	GPD E	160	$2 \cdot 10^7$	10	μ^\pm	NH_3^\uparrow	2022 2 years
Input for Dark Matter Search	\bar{p} production cross section	20-280	$5 \cdot 10^5$	25	p	LH2, LHe	2022 1 month
\bar{p} -induced spectroscopy	Heavy quark exotics	12, 20	$5 \cdot 10^7$	25	\bar{p}	LH2	2022 2 years
Drell-Yan	Pion PDFs	190	$7 \cdot 10^7$	25	π^\pm	C/W	2022 1-2 years

- Expect new Drell-Yan and J/Ψ production data with pion (kaon) beams in the near future !

Summary

- Parton distributions of mesons represent
 - * an interesting topic for theories and experiments
 - * unique opportunities at AMBER, JLab, JPARC and EIC
- J/Ψ production provides useful information on the quark and gluon contents of mesons
 - * Existing data should be included in the global fits for better constraining the gluon distributions in pion and kaon
 - * First results on the extraction of meson PDFs in the framework of statistical model have been obtained using both the Drell-Yan and the J/Ψ data
 - * It would be very interesting to extend the study using other approaches for the global fits.