

# Higgs Potential 2024 (Hefei)

Ref: **JHEP 12 (2023) 018,**

**JHEP 01 (2024) 051,**

***Phys.Lett.B* 833 (2022) 137301**

***arXiv*: 2307.02187**

**Wenxing Zhang(Hebei U.)**

---

## Testing Phase Transition and cosmological history at colliders


---

Collaborators: Yizhou Cai, Hao-Lin Li, Kun Liu, Michael Ramsey-Musolf, Lei Zhang

---

# Bayon asymmetry of the Universe

Cosmic Energy Budget




# Bayon asymmetry of the Universe

## 44 different ways to creat baryons in the Universe

1. GUT baryogenesis
2. GUT baryogenesis after preheating
3. Baryogenesis from primordial black holes
4. String scale baryogenesis
5. Affleck-Dine (AD) baryogenesis
6. Hybridized AD baryogenesis
7. No-scale AD baryogenesis
8. Single field baryogenesis
9. Electroweak (EW) baryogenesis
10. Local EW baryogenesis
11. Non-local EW baryogenesis
12. EW baryogenesis at preheating
13. SUSY EW baryogenesis
14. String mediated EW baryogenesis
15. Baryogenesis via leptogenesis
16. Inflationary baryogenesis
17. Resonant leptogenesis
18. Spontaneous baryogenesis
19. Coherent baryogenesis
20. Gravitational baryogenesis
21. Defect mediated baryogenesis
22. Baryogenesis from long cosmic strings
23. Baryogenesis from short cosmic strings
24. Baryogenesis from collapsing loops

Shaposhnikov, DISCRETE 08, 11, Dec

Cosmic Energy Budget




# Bayon asymmetry of the Universe

## 44 different ways to creat baryons in the Universe

25. Baryogenesis through collapse of vortons
26. Baryogenesis through axion domain walls
27. Baryogenesis through QCD domain walls
28. Baryogenesis through unstable domain walls
29. Baryogenesis from classical force
30. Baryogenesis from electrogenesis
31. B-ball baryogenesis
32. Baryogenesis from CPT breaking
33. Baryogenesis through quantum gravity
34. Baryogenesis via neutrino oscillations
35. Monopole baryogenesis
36. Axino induced baryogenesis
37. Gravitino induced baryogenesis
38. Radion induced baryogenesis
39. Baryogenesis in large extra dimensions
40. Baryogenesis by brane collision
41. Baryogenesis via density fluctuations
42. Baryogenesis from hadronic jets
43. Thermal leptogenesis
44. Nonthermal leptogenesis

Shaposhnikov, DISCRETE 08, 11, Dec

Cosmic Energy Budget



---

# Electroweak Baryogenesis

Sakharov conditions:


- Baryon number violating interactions.
- C and CP violation.
- Departure from thermal equilibrium.

---


# Electroweak Baryogenesis

Sakharov conditions:

- Baryon number violating interactions.
- C and CP violation.
- Departure from thermal equilibrium.



# Electroweak Baryogenesis




Sakharov conditions:

- Baryon number violating interactions.
- C and CP violation.
- Departure from thermal equilibrium.

NEW J.PHYS. 14 (2012) 125003,  
DAVID E.MORRISSEY, MICHAEL J. RAMSEY-MUSOLF

# Electroweak Baryogenesis



Sakharov conditions:

- Baryon number violating interactions.
- C and CP violation.
- Departure from thermal equilibrium.


$$\Gamma_{sph} \sim A_{sph}(T)e^{-E_{sph}(T)/T} > H$$

NEW J.PHYS. 14 (2012) 125003,

DAVID E.MORRISSEY, MICHAEL J. RAMSEY-MUSOLF



# Electroweak Baryogenesis




Sakharov conditions:

- Baryon number violating interactions.
- C and CP violation.
- Departure from thermal equilibrium.

$$\Gamma_{sph} \sim A_{sph}(T)e^{-E_{sph}(T)/T} > H$$

NEW J.PHYS. 14 (2012) 125003,  
DAVID E.MORRISSEY, MICHAEL J. RAMSEY-MUSOLF

# Electroweak Baryogenesis



Sakharov conditions:


- Baryon number violating interactions.
- C and CP violation.
- Departure from thermal equilibrium.

$$\Gamma_{sph} \sim A_{sph}(T)e^{-E_{sph}(T)/T} > H$$

$$S = \frac{n_B(\Delta t_W)}{n_B(0)} > e^{-X}$$

NEW J.PHYS. 14 (2012) 125003,  
DAVID E.MORRISSEY, MICHAEL J. RAMSEY-MUSOLF

# Electroweak Baryogenesis



Sakharov conditions:


- Baryon number violating interactions. ✓
- C and CP violation. ✗
- Departure from thermal equilibrium. ✗

$$\Gamma_{sph} \sim A_{sph}(T)e^{-E_{sph}(T)/T} > H$$

$$S = \frac{n_B(\Delta t_W)}{n_B(0)} > e^{-X}$$

NEW J.PHYS. 14 (2012) 125003,  
DAVID E.MORRISSEY, MICHAEL J. RAMSEY-MUSOLF

# Electroweak Baryogenesis



Sakharov conditions:

- Baryon number violating interactions.
- C and CP violation.
- Departure from thermal equilibrium.

$$\Gamma_{sph} \sim A_{sph}(T)e^{-E_{sph}(T)/T} > H$$

$$S = \frac{n_B(\Delta t_W)}{n_B(0)} > e^{-X}$$

$$\frac{\bar{v}_c(T)}{T} > \zeta_{sph} \sim 1$$

NEW J.PHYS. 14 (2012) 125003,  
DAVID E.MORRISSEY, MICHAEL J. RAMSEY-MUSOLF


---

# SFOEWPT in BSM: Multi-step EWPT

$$V_0(H, S) = -\mu^2(H^\dagger H) + \lambda(H^\dagger H)^2 + \frac{a_1}{2}(H^\dagger H)S + \frac{a_2}{2}(H^\dagger H)S^2 \\ + \frac{b_2}{2}S^2 + \frac{b_3}{3}S^3 + \frac{b_4}{4}S^4,$$

# SFOEWPT in BSM: Multi-step EWPT


$$V_0(H, S) = -\mu^2(H^\dagger H) + \lambda(H^\dagger H)^2 + \frac{a_1}{2}(H^\dagger H)S + \frac{a_2}{2}(H^\dagger H)S^2 + \frac{b_2}{2}S^2 + \frac{b_3}{3}S^3 + \frac{b_4}{4}S^4,$$




EWPT in SM

# SFOEWPT in BSM: Multi-step EWPT

$$V_0(H, S) = -\mu^2(H^\dagger H) + \lambda(H^\dagger H)^2 + \frac{a_1}{2}(H^\dagger H)S + \frac{a_2}{2}(H^\dagger H)S^2 + \frac{b_2}{2}S^2 + \frac{b_3}{3}S^3 + \frac{b_4}{4}S^4,$$




EWPT in SM




1-step EWPT in BSM

# SFOEWPT in BSM: Multi-step EWPT


$$V_0(H, S) = -\mu^2(H^\dagger H) + \lambda(H^\dagger H)^2 + \frac{a_1}{2}(H^\dagger H)S + \frac{a_2}{2}(H^\dagger H)S^2 + \frac{b_2}{2}S^2 + \frac{b_3}{3}S^3 + \frac{b_4}{4}S^4,$$



EWPT in SM



1-step EWPT in BSM



2-step EWPT in BSM




---

# EWPT with a DM candidate: the cxSM

$$\begin{aligned} V_0(H, S) &= \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2 \\ &+ \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 \\ &+ a_1 S + \frac{b_1}{4}S^2 + h.c.. \end{aligned}$$


# EWPT with a DM candidate: the cxSM

$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2$$
$$+ \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4$$
$$+ a_1 S + \frac{b_1}{4}S^2 + h.c..$$



# EWPT with a DM candidate: the cxSM

$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2$$
$$+ \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 \quad U(1)$$
$$+ a_1 S + \frac{b_1}{4}S^2 + h.c..$$




# EWPT with a DM candidate: the cxSM

$$S = x_0 + s + iA$$

$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2 + \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 + a_1S + \frac{b_1}{4}S^2 + h.c..$$


$U(1)$



# EWPT with a DM candidate: the cxSM

$$S = x_0 + s + iA$$

$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2$$
$$+ \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 \quad U(1)$$
$$+ a_1 S + \frac{b_1}{4}S^2 + h.c.. \quad Z_2$$




# EWPT with a DM candidate: the cxSM

$$S = x_0 + s + iA$$

$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2$$
$$+ \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 \quad U(1)$$
$$+ a_1 S + \frac{b_1}{4}S^2 + h.c.. \quad \mathbb{Z}_2$$

AVOID DOMAIN WALL PROBLEM  
EWPT STRENGTH



# EWPT with a DM candidate: the cxSM


$$S = x_0 + s + iA$$

U(1) SSB: MASSIVE PSEUDO-GOLDSTONE BOSON

$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2 + \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 + a_1 S + \frac{b_1}{4}S^2 + h.c..$$

$U(1)$   
 $Z_2$

AVOID DOMAIN WALL PROBLEM  
EWPT STRENGTH



# EWPT with a DM candidate: the cxSM

$$S = x_0 + s + iA$$

U(1) SSB: MASSIVE PSEUDO-GOLDSTONE BOSON


$$V_0(H, S) = \frac{\mu^2}{2} (H^\dagger H) + \frac{\lambda}{4} (H^\dagger H)^2 + \frac{\delta_2}{2} H^\dagger H |S|^2 + \frac{b_2}{2} |S|^2 + \frac{d_2}{4} |S|^4 + a_1 S + \frac{b_1}{4} S^2 + h.c..$$

$\mathcal{M}_h^2 \equiv \begin{pmatrix} M_{hh} & M_{hs} & M_{hA} \\ M_{sh} & M_{ss} & M_{sA} \\ M_{Ah} & M_{As} & M_{AA} \end{pmatrix} = \begin{pmatrix} \frac{1}{2}\lambda v_0^2 & \frac{\delta_2}{2} v_0 v_s & 0 \\ \frac{\delta_2}{2} v_0 v_s & \frac{1}{2} d_2 v_s^2 - \frac{\sqrt{2} a_1}{v_s} & 0 \\ 0 & 0 & -\frac{\sqrt{2} a_1}{v_s} - b_1 \end{pmatrix}$

U(1)

Z<sub>2</sub>

AVOID DOMAIN WALL PROBLEM  
EWPT STRENGTH





# EWPT with a DM candidate: the cxSM

$$S = x_0 + s + iA$$

U(1) SSB: MASSIVE PSEUDO-GOLDSTONE BOSON


$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2 + \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 + a_1 S + \frac{b_1}{4}S^2 + h.c..$$

$\mathcal{M}_h^2 \equiv \begin{pmatrix} M_{hh} & M_{hs} & M_{hA} \\ M_{sh} & M_{ss} & M_{sA} \\ M_{Ah} & M_{As} & M_{AA} \end{pmatrix} = \begin{pmatrix} \frac{1}{2}\lambda v_0^2 & \frac{\delta_2}{2}v_0 v_s & 0 \\ \frac{\delta_2}{2}v_0 v_s & \frac{1}{2}d_2 v_s^2 - \frac{\sqrt{2}a_1}{v_s} & 0 \\ 0 & 0 & -\frac{\sqrt{2}a_1}{v_s} - b_1 \end{pmatrix}$

U(1)

Z<sub>2</sub>

AVOID DOMAIN WALL PROBLEM  
EWPT STRENGTH



# EWPT with a DM candidate: the cxSM

$$S = x_0 + s + iA$$

U(1) SSB: MASSIVE PSEUDO-GOLDSTONE BOSON

$$V_0(H, S) = \frac{\mu^2}{2} (H^\dagger H) + \frac{\lambda}{4} (H^\dagger H)^2 + \frac{\delta_2}{2} H^\dagger H |S|^2 + \frac{b_2}{2} |S|^2 + \frac{d_2}{4} |S|^4 + a_1 S + \frac{b_1}{4} S^2 + h.c..$$



U(1)

Z<sub>2</sub>

AVOID DOMAIN WALL PROBLEM  
EWPT STRENGTH

MIXING & EWPT

$$\mathcal{M}_h^2 \equiv \begin{pmatrix} M_{hh} & M_{hs} & M_{hA} \\ M_{sh} & M_{ss} & M_{sA} \\ M_{Ah} & M_{As} & M_{AA} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \lambda v_0^2 & \frac{\delta_2}{2} v_0 v_s & 0 \\ \frac{\delta_2}{2} v_0 v_s & \frac{1}{2} d_2 v_s^2 - \frac{\sqrt{2} a_1}{v_s} & 0 \\ 0 & 0 & -\frac{\sqrt{2} a_1}{v_s} - b_1 \end{pmatrix}$$



# EWPT with a DM candidate: the cxSM

$$S = x_0 + s + iA$$

U(1) SSB: MASSIVE PSEUDO-GOLDSTONE BOSON

$$V_0(H, S) = \frac{\mu^2}{2} (H^\dagger H) + \frac{\lambda}{4} (H^\dagger H)^2 + \frac{\delta_2}{2} H^\dagger H |S|^2 + \frac{b_2}{2} |S|^2 + \frac{d_2}{4} |S|^4 + a_1 S + \frac{b_1}{4} S^2 + h.c..$$


U(1)

Z<sub>2</sub>

AVOID DOMAIN WALL PROBLEM  
EWPT STRENGTH

MIXING & EWPT

$$\mathcal{M}_h^2 \equiv \begin{pmatrix} M_{hh} & M_{hs} & M_{hA} \\ M_{sh} & M_{ss} & M_{sA} \\ M_{Ah} & M_{As} & M_{AA} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \lambda v_0^2 & \frac{\delta_2}{2} v_0 v_s & 0 \\ \frac{\delta_2}{2} v_0 v_s & \frac{1}{2} d_2 v_s^2 - \frac{\sqrt{2} a_1}{v_s} & 0 \\ 0 & 0 & -\frac{\sqrt{2} a_1}{v_s} - b_1 \end{pmatrix}$$



Theoretical Constraints : Vacuum Stability & Perturbation

# EWPT with a DM candidate: the cxSM

$$S = x_0 + s + iA$$

U(1) SSB: MASSIVE PSEUDO-GOLDSTONE BOSON

$$V_0(H, S) = \frac{\mu^2}{2} (H^\dagger H) + \frac{\lambda}{4} (H^\dagger H)^2 + \frac{\delta_2}{2} H^\dagger H |S|^2 + \frac{b_2}{2} |S|^2 + \frac{d_2}{4} |S|^4 + a_1 S + \frac{b_1}{4} S^2 + h.c..$$

U(1)

Z<sub>2</sub>


AVOID DOMAIN WALL PROBLEM  
EWPT STRENGTH

QUARTIC COUPLINGS & STRENGTH OF EWPT

Theoretical Constraints : Vacuum Stability & Perturbation

MIXING & EWPT

$$\mathcal{M}_h^2 \equiv \begin{pmatrix} M_{hh} & M_{hs} & M_{hA} \\ M_{sh} & M_{ss} & M_{sA} \\ M_{Ah} & M_{As} & M_{AA} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \lambda v_0^2 & \frac{\delta_2}{2} v_0 v_s & 0 \\ \frac{\delta_2}{2} v_0 v_s & \frac{1}{2} d_2 v_s^2 - \frac{\sqrt{2} a_1}{v_s} & 0 \\ 0 & 0 & -\frac{\sqrt{2} a_1}{v_s} - b_1 \end{pmatrix}$$



# EWPT with a DM candidate: the cxSM

$$S = x_0 + s + iA$$

U(1) SSB: MASSIVE PSEUDO-GOLDSTONE BOSON


$$V_0(H, S) = \frac{\mu^2}{2} (H^\dagger H) + \frac{\lambda}{4} (H^\dagger H)^2 + \frac{\delta_2}{2} H^\dagger H |S|^2 + \frac{b_2}{2} |S|^2 + \frac{d_2}{4} |S|^4 + a_1 S + \frac{b_1}{4} S^2 + h.c..$$

$\mathcal{M}_h^2 \equiv \begin{pmatrix} M_{hh} & M_{hs} & M_{hA} \\ M_{sh} & M_{ss} & M_{sA} \\ M_{Ah} & M_{As} & M_{AA} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \lambda v_0^2 & \frac{\delta_2}{2} v_0 v_s & 0 \\ \frac{\delta_2}{2} v_0 v_s & \frac{1}{2} d_2 v_s^2 - \frac{\sqrt{2} a_1}{v_s} & 0 \\ 0 & 0 & -\frac{\sqrt{2} a_1}{v_s} - b_1 \end{pmatrix}$

MIXING & EWPT

AVOID DOMAIN WALL PROBLEM  
EWPT STRENGTH

QUARTIC COUPLINGS & STRENGTH OF EWPT



**Theoretical Constraints : Vacuum Stability & Perturbation**

**Experimental Constraints : EW Precision Observables & Higgs Measurement**

# EWPT with a DM candidate: the cxSM

$$S = x_0 + s + iA$$

U(1) SSB: MASSIVE PSEUDO-GOLDSTONE BOSON

$$V_0(H, S) = \frac{\mu^2}{2} (H^\dagger H) + \frac{\lambda}{4} (H^\dagger H)^2 + \frac{\delta_2}{2} H^\dagger H |S|^2 + \frac{b_2}{2} |S|^2 + \frac{d_2}{4} |S|^4 + a_1 S + \frac{b_1}{4} S^2 + h.c..$$

$\mathcal{M}_h^2 \equiv \begin{pmatrix} M_{hh} & M_{hs} & M_{hA} \\ M_{sh} & M_{ss} & M_{sA} \\ M_{Ah} & M_{As} & M_{AA} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \lambda v_0^2 & \frac{\delta_2}{2} v_0 v_s & 0 \\ \frac{\delta_2}{2} v_0 v_s & \frac{1}{2} d_2 v_s^2 - \frac{\sqrt{2} a_1}{v_s} & 0 \\ 0 & 0 & -\frac{\sqrt{2} a_1}{v_s} - b_1 \end{pmatrix}$


MIXING & EWPT

U(1)

Z<sub>2</sub>

AVOID DOMAIN WALL PROBLEM  
EWPT STRENGTH

QUARTIC COUPLINGS & STRENGTH OF EWPT



Theoretical Constraints : Vacuum Stability & Perturbation

Experimental Constraints : EW Precision Observables & Higgs Measurement

MIXING ANGLE & DM & HEAVY HIGGS MASS & EWPT & COLLIDER

---

# Constraints: $W$ mass measurement

$$\Delta\mathcal{O} = (\cos^2\theta - 1)\mathcal{O}^{\text{SM}}(m_{h_1}) + \sin^2\theta\mathcal{O}^{\text{SM}}(m_{h_2}) = \sin^2\theta [\mathcal{O}^{\text{SM}}(m_{h_2}) - \mathcal{O}^{\text{SM}}(m_{h_1})]$$

$$\Delta S = 0.086 \pm 0.077,$$

$$\Delta T = 0.177 \pm 0.070$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.89 \\ 0.89 & 1 \end{pmatrix}.$$

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$

# Constraints: W mass measurement

$$\Delta\mathcal{O} = (\cos^2\theta - 1)\mathcal{O}^{\text{SM}}(m_{h_1}) + \sin^2\theta\mathcal{O}^{\text{SM}}(m_{h_2}) = \sin^2\theta [\mathcal{O}^{\text{SM}}(m_{h_2}) - \mathcal{O}^{\text{SM}}(m_{h_1})]$$

$$\Delta S = 0.086 \pm 0.077,$$

$$\Delta T = 0.177 \pm 0.070$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.89 \\ 0.89 & 1 \end{pmatrix}.$$


Two degree of freedom, 95% C.L.

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$



# Constraints: W mass measurement

$$\Delta\mathcal{O} = (\cos^2\theta - 1)\mathcal{O}^{\text{SM}}(m_{h_1}) + \sin^2\theta\mathcal{O}^{\text{SM}}(m_{h_2}) = \sin^2\theta [\mathcal{O}^{\text{SM}}(m_{h_2}) - \mathcal{O}^{\text{SM}}(m_{h_1})]$$



$$\Delta S = 0.086 \pm 0.077,$$

$$\Delta T = 0.177 \pm 0.070$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.89 \\ 0.89 & 1 \end{pmatrix}.$$


Two degree of freedom, 95% C.L.

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$

TRANSLATE FROM CDFII W-MASS MEASUREMENT

# Constraints: W mass measurement

$$\Delta\mathcal{O} = (\cos^2\theta - 1)\mathcal{O}^{\text{SM}}(m_{h_1}) + \sin^2\theta\mathcal{O}^{\text{SM}}(m_{h_2}) = \sin^2\theta [\mathcal{O}^{\text{SM}}(m_{h_2}) - \mathcal{O}^{\text{SM}}(m_{h_1})]$$



Small  $m_{h_2}$  with large mixing angle is favored!

$$\Delta S = 0.086 \pm 0.077,$$

$$\Delta T = 0.177 \pm 0.070$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.89 \\ 0.89 & 1 \end{pmatrix}.$$


Two degree of freedom, 95% C.L.

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$


TRANSLATE FROM CDFII W-MASS MEASUREMENT

# Constraints: W mass measurement

$$\Delta\mathcal{O} = (\cos^2\theta - 1)\mathcal{O}^{\text{SM}}(m_{h_1}) + \sin^2\theta\mathcal{O}^{\text{SM}}(m_{h_2}) = \sin^2\theta [\mathcal{O}^{\text{SM}}(m_{h_2}) - \mathcal{O}^{\text{SM}}(m_{h_1})]$$



Small  $m_{h_2}$  with large mixing angle is favored!



$$\Delta S = 0.086 \pm 0.077,$$

$$\Delta T = 0.177 \pm 0.070$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.89 \\ 0.89 & 1 \end{pmatrix}.$$


Two degree of freedom, 95% C.L.

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$


TRANSLATE FROM CDFII W-MASS MEASUREMENT

# Constraints: W mass measurement

$$\Delta\mathcal{O} = (\cos^2\theta - 1)\mathcal{O}^{\text{SM}}(m_{h_1}) + \sin^2\theta\mathcal{O}^{\text{SM}}(m_{h_2}) = \sin^2\theta [\mathcal{O}^{\text{SM}}(m_{h_2}) - \mathcal{O}^{\text{SM}}(m_{h_1})]$$



Small  $m_{h_2}$  with large mixing angle is favored!



$$\Delta S = 0.086 \pm 0.077,$$

$$\Delta T = 0.177 \pm 0.070$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.89 \\ 0.89 & 1 \end{pmatrix}.$$

Two degree of freedom, 95% C.L.


$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$

TRANSLATE FROM CDFII W-MASS MEASUREMENT


DOES THE W MASS MEASUREMENT EXCLUDE THE SINGLET HIGGS MODEL ?

# Constraints: W mass measurement

$$\Delta\mathcal{O} = (\cos^2\theta - 1)\mathcal{O}^{\text{SM}}(m_{h_1}) + \sin^2\theta\mathcal{O}^{\text{SM}}(m_{h_2}) = \sin^2\theta [\mathcal{O}^{\text{SM}}(m_{h_2}) - \mathcal{O}^{\text{SM}}(m_{h_1})]$$



Small  $m_{h_2}$  with large mixing angle is favored!



$$\Delta S = 0.086 \pm 0.077,$$

$$\Delta T = 0.177 \pm 0.070$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.89 \\ 0.89 & 1 \end{pmatrix}.$$

Two degree of freedom, 95% C.L.

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$

TRANSLATE FROM CDFII W-MASS MEASUREMENT

DOES THE W MASS MEASUREMENT EXCLUDE THE SINGLET HIGGS MODEL ?

**The answer is NO!!**

This is not a UV complete model

# Conservative constraints : EWPO & Higgs Measurement

*Eur.Phys.J.C* 78 (2018) 8

$$S - S_{SM} = 0.04 \pm 0.11$$

$$T - T_{SM} = 0.09 \pm 0.14$$

$$U - U_{SM} = -0.02 \pm 0.11$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.92 & -0.68 \\ 0.92 & 1 & -0.87 \\ -0.68 & -0.87 & 1 \end{pmatrix}.$$

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$

# Conservative constraints : EWPO & Higgs Measurement

*Eur.Phys.J.C* 78 (2018) 8


$$S - S_{SM} = 0.04 \pm 0.11$$

$$T - T_{SM} = 0.09 \pm 0.14$$

$$U - U_{SM} = -0.02 \pm 0.11$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.92 & -0.68 \\ 0.92 & 1 & -0.87 \\ -0.68 & -0.87 & 1 \end{pmatrix}.$$

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$



*JHEP* 01 (2024) 051,  
WX.Z, Y.Cai, M.J.Ramsey-Musolf, L.Zhang

# Conservative constraints : EWPO & Higgs Measurement

*Eur.Phys.J.C* 78 (2018) 8


$$S - S_{SM} = 0.04 \pm 0.11$$

$$T - T_{SM} = 0.09 \pm 0.14$$

$$U - U_{SM} = -0.02 \pm 0.11$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.92 & -0.68 \\ 0.92 & 1 & -0.87 \\ -0.68 & -0.87 & 1 \end{pmatrix}.$$

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$



*JHEP* 01 (2024) 051,  
WX.Z, Y.Cai, M.J.Ramsey-Musolf, L.Zhang

New physics may induce deviation in Higgs couplings. Therefore it modifies the Higgs signal strength in Higgs measurement.

Production mode	ggF+ $b\bar{b}H$	VBF	WH	ZH	$t\bar{t}H$	$tH$
$\sum_f \mu_{i \rightarrow h_1 \rightarrow ff}$	$1.03^{+0.07}_{-0.07}$	$1.10^{+0.13}_{-0.12}$	$1.16^{+0.23}_{-0.22}$	$0.96^{+0.22}_{-0.21}$	$0.74^{+0.24}_{-0.24}$	$6.61^{+4.24}_{-3.76}$

*Nature* 607, 52-59 (2022)

$$\mu_{pp \rightarrow h_1 \rightarrow XX} = \frac{\sigma_{pp \rightarrow h_1} BR(h_1 \rightarrow XX)}{\sigma_{pp \rightarrow h}^{SM} BR(h \rightarrow XX)_{SM}} \simeq \cos^2 \theta,$$

$$\chi^2 = \sum_{i,f} \frac{(\mu_{i \rightarrow h_1 \rightarrow f}^{xSM} - \mu_{i \rightarrow h_1 \rightarrow f}^{obs})^2}{\sigma_{\mu_{i \rightarrow h \rightarrow f}}^2}, \quad \Delta\chi^2 = \chi^2 - \chi_{min}^2 < 3.841. \rightarrow \text{For 1 DoF, 95\% C.L.}$$

This set  $\|\sin \theta\| < 0.2$ .



# Conservative constraints : EWPO & Higgs Measurement

*Eur.Phys.J.C* 78 (2018) 8


$$S - S_{SM} = 0.04 \pm 0.11$$

$$T - T_{SM} = 0.09 \pm 0.14$$

$$U - U_{SM} = -0.02 \pm 0.11$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.92 & -0.68 \\ 0.92 & 1 & -0.87 \\ -0.68 & -0.87 & 1 \end{pmatrix}.$$

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$



*JHEP* 01 (2024) 051,  
WX.Z, Y.Cai, M.J.Ramsey-Musolf, L.Zhang

New physics may induce deviation in Higgs couplings. Therefore it modifies the Higgs signal strength in Higgs measurement.

Production mode	ggF+ $b\bar{b}H$	VBF	WH	ZH	$t\bar{t}H$	$tH$
$\sum_f \mu_{i \rightarrow h_1 \rightarrow ff}$	$1.03^{+0.07}_{-0.07}$	$1.10^{+0.13}_{-0.12}$	$1.16^{+0.23}_{-0.22}$	$0.96^{+0.22}_{-0.21}$	$0.74^{+0.24}_{-0.24}$	$6.61^{+4.24}_{-3.76}$

*Nature* 607, 52-59 (2022)

$$\mu_{pp \rightarrow h_1 \rightarrow XX} = \frac{\sigma_{pp \rightarrow h_1} BR(h_1 \rightarrow XX)}{\sigma_{pp \rightarrow h}^{SM} BR(h \rightarrow XX)_{SM}} \simeq \cos^2 \theta,$$

$$\chi^2 = \sum_{i,f} \frac{(\mu_{i \rightarrow h_1 \rightarrow f}^{xSM} - \mu_{i \rightarrow h_1 \rightarrow f}^{obs})^2}{\sigma_{\mu_{i \rightarrow h \rightarrow f}}^2}, \quad \Delta\chi^2 = \chi^2 - \chi_{min}^2 < 3.841. \rightarrow \text{For 1 DoF, 95\% C.L.}$$

**This set  $\|\sin \theta\| < 0.2$ .**

# Conservative constraints : EWPO & Higgs Measurement

*Eur.Phys.J.C* 78 (2018) 8


$$S - S_{SM} = 0.04 \pm 0.11$$

$$T - T_{SM} = 0.09 \pm 0.14$$

$$U - U_{SM} = -0.02 \pm 0.11$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.92 & -0.68 \\ 0.92 & 1 & -0.87 \\ -0.68 & -0.87 & 1 \end{pmatrix}.$$

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$



*JHEP* 01 (2024) 051,  
WX.Z, Y.Cai, M.J.Ramsey-Musolf, L.Zhang

New physics may induce deviation in Higgs couplings. Therefore it modifies the Higgs signal strength in Higgs measurement.

Production mode	ggF+ $b\bar{b}H$	VBF	WH	ZH	$t\bar{t}H$	$tH$
$\sum_f \mu_{i \rightarrow h_1 \rightarrow ff}$	$1.03^{+0.07}_{-0.07}$	$1.10^{+0.13}_{-0.12}$	$1.16^{+0.23}_{-0.22}$	$0.96^{+0.22}_{-0.21}$	$0.74^{+0.24}_{-0.24}$	$6.61^{+4.24}_{-3.76}$

*Nature* 607, 52-59 (2022)

$$\mu_{pp \rightarrow h_1 \rightarrow XX} = \frac{\sigma_{pp \rightarrow h_1} BR(h_1 \rightarrow XX)}{\sigma_{pp \rightarrow h}^{SM} BR(h \rightarrow XX)_{SM}} \simeq \cos^2 \theta,$$

$$\chi^2 = \sum_{i,f} \frac{(\mu_{i \rightarrow h_1 \rightarrow f}^{xSM} - \mu_{i \rightarrow h_1 \rightarrow f}^{obs})^2}{\sigma_{\mu_{i \rightarrow h \rightarrow f}}^2}, \quad \Delta\chi^2 = \chi^2 - \chi_{min}^2 < 3.841. \rightarrow \text{For 1 DoF, 95\% C.L.}$$

This set  $\|\sin \theta\| < 0.2$ .

# Conservative constraints : EWPO & Higgs Measurement

*Eur.Phys.J.C* 78 (2018) 8


$$S - S_{SM} = 0.04 \pm 0.11$$

$$T - T_{SM} = 0.09 \pm 0.14$$

$$U - U_{SM} = -0.02 \pm 0.11$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.92 & -0.68 \\ 0.92 & 1 & -0.87 \\ -0.68 & -0.87 & 1 \end{pmatrix}.$$

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$



*JHEP* 01 (2024) 051,  
WX.Z, Y.Cai, M.J.Ramsey-Musolf, L.Zhang

New physics may induce deviation in Higgs couplings. Therefore it modifies the Higgs signal strength in Higgs measurement.

Production mode	ggF+ $b\bar{b}H$	VBF	WH	ZH	$t\bar{t}H$	$tH$
$\sum_f \mu_{i \rightarrow h_1 \rightarrow ff}$	$1.03^{+0.07}_{-0.07}$	$1.10^{+0.13}_{-0.12}$	$1.16^{+0.23}_{-0.22}$	$0.96^{+0.22}_{-0.21}$	$0.74^{+0.24}_{-0.24}$	$6.61^{+4.24}_{-3.76}$

*Nature* 607, 52-59 (2022)

$$\mu_{pp \rightarrow h_1 \rightarrow XX} = \frac{\sigma_{pp \rightarrow h_1} BR(h_1 \rightarrow XX)}{\sigma_{pp \rightarrow h}^{SM} BR(h \rightarrow XX)_{SM}} \simeq \cos^2 \theta,$$

$$\chi^2 = \sum_{i,f} \frac{(\mu_{i \rightarrow h_1 \rightarrow f}^{xSM} - \mu_{i \rightarrow h_1 \rightarrow f}^{obs})^2}{\sigma_{\mu_{i \rightarrow h \rightarrow f}}^2}, \quad \Delta\chi^2 = \chi^2 - \chi_{min}^2 < 3.841. \rightarrow \text{For 1 DoF, 95\% C.L.}$$

This set  $\|\sin \theta\| < 0.2$ .

# Conservative constraints : EWPO & Higgs Measurement

*Eur.Phys.J.C* 78 (2018) 8


$$S - S_{SM} = 0.04 \pm 0.11$$

$$T - T_{SM} = 0.09 \pm 0.14$$

$$U - U_{SM} = -0.02 \pm 0.11$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.92 & -0.68 \\ 0.92 & 1 & -0.87 \\ -0.68 & -0.87 & 1 \end{pmatrix}.$$

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$



*JHEP* 01 (2024) 051,  
WX.Z, Y.Cai, M.J.Ramsey-Musolf, L.Zhang

New physics may induce deviation in Higgs couplings. Therefore it modifies the Higgs signal strength in Higgs measurement.

Production mode	ggF+ $b\bar{b}H$	VBF	WH	ZH	$t\bar{t}H$	$tH$
$\sum_f \mu_{i \rightarrow h_1 \rightarrow ff}$	$1.03^{+0.07}_{-0.07}$	$1.10^{+0.13}_{-0.12}$	$1.16^{+0.23}_{-0.22}$	$0.96^{+0.22}_{-0.21}$	$0.74^{+0.24}_{-0.24}$	$6.61^{+4.24}_{-3.76}$

**B-jet**

Nature 607, 52-59 (2022)

$$\mu_{pp \rightarrow h_1 \rightarrow XX} = \frac{\sigma_{pp \rightarrow h_1} BR(h_1 \rightarrow XX)}{\sigma_{pp \rightarrow h}^{SM} BR(h \rightarrow XX)_{SM}} \simeq \cos^2 \theta,$$

$$\chi^2 = \sum_{i,f} \frac{(\mu_{i \rightarrow h_1 \rightarrow f}^{xSM} - \mu_{i \rightarrow h_1 \rightarrow f}^{obs})^2}{\sigma_{\mu_{i \rightarrow h \rightarrow f}}^2}, \quad \Delta\chi^2 = \chi^2 - \chi_{min}^2 < 3.841. \rightarrow \text{For 1 DoF, 95\% C.L.}$$

This set  $\|\sin \theta\| < 0.2$ .

**HIGGS MEASUREMENT AT THE LHC UTILIZES B-JET DECAY MODES MOSTLY.  
HIGGS DECAY TO LIGHT QUARKS WOULD IMPROVE THE HIGGS MEASUREMENT SIGNIFICANTLY**

# Conservative constraints : EWPO & Higgs Measurement

Eur.Phys.J.C 78 (2018) 8


$$S - S_{SM} = 0.04 \pm 0.11$$

$$T - T_{SM} = 0.09 \pm 0.14$$

$$U - U_{SM} = -0.02 \pm 0.11$$

$$\rho_{ij} = \begin{pmatrix} 1 & 0.92 & -0.68 \\ 0.92 & 1 & -0.87 \\ -0.68 & -0.87 & 1 \end{pmatrix}.$$

$$\chi^2 = (X - \hat{X})_i (\sigma^2)_{ij}^{-1} (X - \hat{X})_j < 5.99$$



JHEP 01 (2024) 051,  
WX.Z, Y.Cai, M.J.Ramsey-Musolf, L.Zhang

New physics may induce deviation in Higgs couplings. Therefore it modifies the Higgs signal strength in Higgs measurement.

Production mode	ggF+ $b\bar{b}H$	VBF	WH	ZH	$t\bar{t}H$	$tH$
$\sum_f \mu_{i \rightarrow h_1 \rightarrow ff}$	$1.03^{+0.07}_{-0.07}$	$1.10^{+0.13}_{-0.12}$	$1.16^{+0.23}_{-0.22}$	$0.96^{+0.22}_{-0.21}$	$0.74^{+0.24}_{-0.24}$	$6.61^{+4.24}_{-3.76}$

**B-jet**

Nature 607, 52-59 (2022)

$$\mu_{pp \rightarrow h_1 \rightarrow XX} = \frac{\sigma_{pp \rightarrow h_1} BR(h_1 \rightarrow XX)}{\sigma_{pp \rightarrow h}^{SM} BR(h \rightarrow XX)_{SM}} \simeq \cos^2 \theta,$$

$$\chi^2 = \sum_{i,f} \frac{(\mu_{i \rightarrow h_1 \rightarrow f}^{xSM} - \mu_{i \rightarrow h_1 \rightarrow f}^{obs})^2}{\sigma_{\mu_{i \rightarrow h \rightarrow f}}^2}, \quad \Delta\chi^2 = \chi^2 - \chi_{min}^2 < 3.841. \rightarrow \text{For 1 DoF, 95\% C.L.}$$

This set  $\|\sin \theta\| < 0.2$ .

**HIGGS MEASUREMENT AT THE LHC UTILIZES B-JET DECAY MODES MOSTLY.  
HIGGS DECAY TO LIGHT QUARKS WOULD IMPROVE THE HIGGS MEASUREMENT SIGNIFICANTLY**

**OPPORTUNITIES AT FUTURE COLLIDERS!!!**

---

# Detect the SFOEWPT at colliders

## TAKE-AWAY MESSAGE:

On one hand, enough strong first order EWPT needs considerable new physics contribution. On the other hand, too big new physics coupling leads to deviation from experiment observation.

---

# Detect the SFOEWPT at colliders

## TAKE-AWAY MESSAGE:

On one hand, enough strong first order EWPT needs considerable new physics contribution. On the other hand, too big new physics coupling leads to deviation from experiment observation.


**The viable SFOEWPT model is possible to be fully tested in the current and future colliders**

# Detect the SFOEWPT at colliders

## TAKE-AWAY MESSAGE:

On one hand, enough strong first order EWPT needs considerable new physics contribution. On the other hand, too big new physics coupling leads to deviation from experiment observation.

The viable SFOEWPT model is possible to be fully tested in the current and future colliders






# Detect the SFOEWPT at colliders

## TAKE-AWAY MESSAGE:

On one hand, enough strong first order EWPT needs considerable new physics contribution. On the other hand, too big new physics coupling leads to deviation from experiment observation.

**The viable SFOEWPT model is possible to be fully tested in the current and future colliders**





**CEPC tests the lower mass region.**

# Detect the SFOEWPT at colliders

## TAKE-AWAY MESSAGE:

On one hand, enough strong first order EWPT needs considerable new physics contribution. On the other hand, too big new physics coupling leads to deviation from experiment observation.

The viable SFOEWPT model is possible to be fully tested in the current and future colliders




CEPC tests the lower mass region.

# Detect the SFOEWPT at colliders


## TAKE-AWAY MESSAGE:

On one hand, enough strong first order EWPT needs considerable new physics contribution. On the other hand, too big new physics coupling leads to deviation from experiment observation.

The viable SFOEWPT model is possible to be fully tested in the current and future colliders



CEPC tests the lower mass region.




LHC tests the higher mass region.

# Detect the SFOEWPT at colliders


**TAKE-AWAY MESSAGE:**

On one hand, enough strong first order EWPT needs considerable new physics contribution. On the other hand, too big new physics coupling leads to deviation from experiment observation.

**The viable SFOEWPT model is possible to be fully tested in the current and future colliders**



**CEPC tests the lower mass region.**



**LHC tests the higher mass region.**


**For both CEPC and the LHC, multiple channels are investigated. We must ask: Has the information from these detection channels been fully utilized?**

# Detect the SFOEWPT at colliders


## TAKE-AWAY MESSAGE:

On one hand, enough strong first order EWPT needs considerable new physics contribution. On the other hand, too big new physics coupling leads to deviation from experiment observation.

The viable SFOEWPT model is possible to be fully tested in the current and future colliders



CEPC tests the lower mass region.



LHC tests the higher mass region.

For both CEPC and the LHC, multiple channels are investigated. We must ask: Has the information from these detection channels been fully utilized?


IF NOT, HOW CAN THESE DETECTION CHANNELS TO BE OPTIMIZED?

# Detect the SFOEWPT at colliders


## TAKE-AWAY MESSAGE:

On one hand, enough strong first order EWPT needs considerable new physics contribution. On the other hand, too big new physics coupling leads to deviation from experiment observation.

The viable SFOEWPT model is possible to be fully tested in the current and future colliders



CEPC tests the lower mass region.








LHC tests the higher mass region.

For both CEPC and the LHC, multiple channels are investigated. We must ask: Has the information from these detection channels been fully utilized?

IF NOT, HOW CAN THESE DETECTION CHANNELS TO BE OPTIMIZED?

Two answers are included in this talk

# Optimizing detecting PT ability at colliders : Combine $b\bar{b}\gamma\gamma$ and $ZZ \rightarrow 4\ell$



**THE DETECTABLE SPACE HAS EXPANDED !!!**

---

# Detect the DM phenomenology and SFOEWPT at colliders

$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2 + \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 \\ + a_1 S + \frac{b_1}{4}S^2 + h.c..$$

Nonzero coupling  $g_{1AA}$  would induce Higgs invisible decay if  $m_A < \frac{m_{h_1}}{2}$

$$g_{1AA} = \frac{\sqrt{2}a_1 + m_{h_1}^2 v_s}{2v_s^2} \sin \theta,$$





# Detect the DM phenomenology and SFOEWPT at colliders

$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2 + \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 \\ + a_1 S + \frac{b_1}{4}S^2 + h.c..$$

Nonzero coupling  $g_{1AA}$  would induce Higgs invisible decay if  $m_A < \frac{m_{h_1}}{2}$

$$g_{1AA} = \frac{\sqrt{2}a_1 + m_{h_1}^2 v_s}{2v_s^2} \sin \theta,$$

The decay width is  $\Gamma_{h_1 \rightarrow AA} \sim \sqrt{1 - \frac{4m_A^2}{m_{h_1}^2}} \times \left(\frac{\sin \theta}{0.1}\right)^2 \times 50$  [MeV]

$$\Gamma_{H \rightarrow inv}^{SM} \lesssim 4.1 \text{ MeV} \times 13\% \simeq 0.4 \text{ MeV}$$

# Detect the DM phenomenology and SFOEWPT at colliders

$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2 + \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 \\ + a_1 S + \frac{b_1}{4}S^2 + h.c..$$

Nonzero coupling  $g_{1AA}$  would induce Higgs invisible decay if  $m_A < \frac{m_{h_1}}{2}$

$$g_{1AA} = \frac{\sqrt{2}a_1 + m_{h_1}^2 v_s}{2v_s^2} \sin \theta,$$

The decay width is  $\Gamma_{h_1 \rightarrow AA} \sim \sqrt{1 - \frac{4m_A^2}{m_{h_1}^2}} \times \left(\frac{\sin \theta}{0.1}\right)^2 \times 50$  [MeV]

$$\Gamma_{H \rightarrow inv}^{SM} \lesssim 4.1 \text{ MeV} \times 13\% \simeq 0.4 \text{ MeV}$$

Thus it is safe to set the DM mass with  $m_A \geq 62.5$  GeV

$m_A > 62.5$  GeV: Recently-studied case

$m_A = 62.5$  GeV: Well-studied case

# Detect the DM phenomenology and SFOEWPT at colliders

$$V_0(H, S) = \frac{\mu^2}{2}(H^\dagger H) + \frac{\lambda}{4}(H^\dagger H)^2 + \frac{\delta_2}{2}H^\dagger H|S|^2 + \frac{b_2}{2}|S|^2 + \frac{d_2}{4}|S|^4 + a_1 S + \frac{b_1}{4}S^2 + h.c..$$

Nonzero coupling  $g_{1AA}$  would induce Higgs invisible decay if  $m_A < \frac{m_{h_1}}{2}$

$$g_{1AA} = \frac{\sqrt{2}a_1 + m_{h_1}^2 v_s}{2v_s^2} \sin \theta,$$




The decay width is  $\Gamma_{h_1 \rightarrow AA} \sim \sqrt{1 - \frac{4m_A^2}{m_{h_1}^2}} \times \left(\frac{\sin \theta}{0.1}\right)^2 \times 50$  [MeV]

$$\Gamma_{H \rightarrow inv}^{SM} \lesssim 4.1 \text{ MeV} \times 13\% \simeq 0.4 \text{ MeV}$$

Thus it is safe to set the DM mass with  $m_A \geq 62.5$  GeV

$m_A > 62.5$  GeV: Recently-studied case

$m_A = 62.5$  GeV: Well-studied case









Recently-studied case

Well-studied case

For  $m_A > 62.5$  GeV, most SFOPT points survive XENON1T search and are able to be detected by XENON-nT or PandaX-4T.

For  $m_A = 62.5$  GeV, most SFOPT points do not survive XENON1T search.


# cxSM: search EWPT & DM at the (HL-)LHC




Di-Boson: Cover the whole space in  $m_{h_2} > 600$  GeV

Di-Higgs: Cover significant portion in  $m_{h_2} < 400$  GeV


LHC Run-II : Almost no exclusion ability



# cxSM: search for $b\bar{b} + \text{MET}$ at the (HL-)LHC



Detectable region by singlet search only,  
including di-boson and di-higgs channels



Space that can only be detected by  $b\bar{b} + \text{MET}$

---

# Summary

We briefly see how the Higgs measurement impact on the EWPT searches at colliders, which gives an opportunity to the future collider—CEPC. ML method is expected to optimize the preselection step.

Two methods are introduced to optimize the heavy Higgs search for EWPT. One is to combine di-Higgs and di-boson channels, the other is to make use the branching new search channel  $b\bar{b} + \text{MET}$ .

Without updating LHC hardware, detectable parameter space are extended by the first method. DM is very hopefully to be discovered by the future Xenon-nT or PandaX-4T.

---