



上海交通大學
SHANGHAI JIAO TONG UNIVERSITY

李政道研究所
Tsung-Dao Lee Institute

Tracking system R&D for the DarkSHINE experiment

Dian YU, on behalf of the DarkSHINE R&D team

Tsung-Dao Lee Institute



Shaan Xi, April 17-20 2025

Physics motivation




上海交通大学
SHANGHAI JIAO TONG UNIVERSITY

李政道研究所
Tsung-Dao Lee Institute




- Cosmological observation has proved the existence of dark matter
- No evidence that dark matter as WIMP exists yet, a large parameter space ruled out in GeV~TeV mass range
- In the sub-GeV range, **dark photon** hypothesis is brought out as the force carrier and portal between ordinary matter and dark sector

Dark photon production modes




VISIBLE DECAY MODE $m'_A < 2m_X$



Having two interaction vertices \rightarrow production rate highly suppressed

$$N \propto \epsilon^4 \ll N \propto \epsilon^2(1 - \epsilon^2) \approx \epsilon^2$$

INVISIBLE DECAY MODE $m'_A > 2m_X$



Focusing on invisible decays of dark photon into light dark matter

- **Experimental signatures:** missing energy, **missing momentum**.
- **DarkSHINE:** single electron on target experiment, searching for dark photon.

Prospective studies (a year = $3e14$ EOT)





上海交通大学
SHANGHAI JIAO TONG UNIVERSITY

李政道研究所
Tsung-Dao Lee Institute

- The DarkSHINE experiment can provide competitive results on hunting for dark photon (left), also very sensitive to some popular dark matter models (right).

Sci. China-Phys. Mech. Astron., 66(1): 211062 (2023); *arxiv:2411.09345*





DarkSHINE experimental approaches

- **High repetition rate single electron beam**


- The beam can be provided by SHINE based on SXFEL
- Properties : 8GeV, 10MHz, single electron on target

- **Energy + Momentum loss detection**

- Tracking system, ECAL, HCAL



Conceptual diagram of the detector




Tracking system design





上海交通大学
SHANGHAI JIAO TONG UNIVERSITY

李政道研究所
Tsung-Dao Lee Institute



[arXiv:2310.13926](https://arxiv.org/abs/2310.13926)

[Nucl Sci Tech 35, 201 \(2024\)](#)



Pitch (μm)	Strip (μm)
100	50
60	40
45	30

- Silicon strips detector under 1.5T magnetic field , $\sim 10 \mu\text{m}$ position resolution.
- 7 layers of tagging + 6 layers of recoil tracker, two silicon strips sensors each layer to enable resolution in y axis.
- AC-LGAD silicon strip sensor $3638 \times 3638 \mu\text{m}^2$ for performance study.
- In collaboration with IHEP-CAS HGTD team





Performance

arXiv:2310.13926
Nucl Sci Tech 35, 201 (2024)





上海交通大学
SHANGHAI JIAO TONG UNIVERSITY



李政道研究所
Tsung-Dao Lee Institute





- Performance testing: laser beam as signal source
spot diameter $6 \sim 10 \mu m$
- Performance test: step $2 \mu m$, 1000+ scans per point




- Spatial resolution
 $7 \sim 12 \mu\text{m}$



- Time performance
 $6 \sim 14\text{ps}$



- IDE1140 is chosen as the preamplifier-shaper ASIC.
- Power board for ASICs and sensor powering is developed. We are still working on **sensor board, amplifier and ADC**.
- The design is referred to AMS experiment, thanks Prof Zijun Xu and Mengke Cai from IHEP




- Design the full electronics
- Manufacture the prototype
- Beamtest, etc.

• **Welcome your collaboration!**

- **DarkSHINE: a fixed-target experiment searching for dark photon invisible decay.**
- **Competitive sensitivity has been studied for dark photon and light dark matter**
 - *Sci. China-Phys. Mech. Astron.*, 66(1): 211062 (2023).
- **Detector key technology R&D updates have been presented**
 - AC-LGAD silicon strips: *Nucl.Sci.Tech.* 35 (2024) 11, 201.
 - LYSO+SiPM ECAL: *Nucl.Sci.Tech.* 36(2025) 3, 41.
 - Scintillator + Iron absorber HCAL: *Nucl.Sci.Tech.* 35 (2024) 9, 148.
- **Aiming for detector prototype manufacture and CDR.**

Backup



where x is the impact position, f_i is the signal fraction of each channel, α_i is the signal fraction of each channel at $x = 0$, and β_i is the change rate of the signal fraction of each channel with the impact position. We set $x = 0$ at centre of

$$f_i = \frac{A_{\max}^i}{A_{\max}^1 + A_{\max}^2 + A_{\max}^3 + A_{\max}^4}, i = 1, 2, 3, 4$$

$$\sigma_{\text{time}} = \sigma(t_1 + t_2 - t_3 - t_4)/4$$


Introduce extra $U(1)_X$ symmetry \rightarrow New Gauge Field $X \rightarrow$ Dark Photon Mediator A'
 $U(1)_{\text{em}} \rightarrow U(1)_{\text{em}} \times U(1)_X$

5th fundamental interaction in our universe

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + A_{\mu}j_{\text{em}}^{\mu} - \frac{1}{4}X_{\mu\nu}X^{\mu\nu} + X_{\mu}j_X^{\mu}$$

SM Photon γ

Dark Photon A'




- A' & γ kin. mixing
- Renormalizable and Gauge Invariant
- Straightforward for experimental search
- Free param, kin. mixing (ϵ), mass ($m_{A'}$)

4 Renormalizable "Portals"

Axion	$\frac{1}{f_a} F_{\mu\nu} \tilde{F}^{\mu\nu} a$	Axion/ALP	Higgs	$\lambda H^2 S^2 + \mu H^2 S$	Exotic Higgs decay?
Vector	$\epsilon F^{Y,\mu\nu} F'_{\mu\nu}$	Dark photon	Neutrino	$\kappa (HL)N$	Sterile neutrino?

Direct Detection

Accelerator based experiments



• In accelerator-based experiments, this difference can be reduced to $10^2 \sim 10^3$ orders of magnitudes, due to the fact of insensitive to DM's mass and spin in its production.