

Searching for apparent baryon number violation in Λ_c^+ decays at STCF

Zeren Simon Wang 王泽人

Hefei University of Technology 合肥工业大学

Work in progress

in collaboration with Xin-Ru Tang, Yu Zhang, Yu Zhang, and Xiaorong Zhou


November 26th, 2025

The 7th International Workshop on Future Tau Charm Facilities
Huangshan, Anhui, China



Motivation

- B : conserved perturbatively in the SM
- B : violated only through non-perturbative effects such as sphalerons and instantons: strongly suppressed at low energies
- Matter-antimatter asymmetry \Rightarrow baryogenesis:
Sakharov conditions including B
- No observation of BNV processes experimentally so far and
Increasingly stringent bounds
- Study neutrino-extended EFTs with BNV and ν_s : ν SMEFT & ν LEFT



- STCF, large $\Lambda_c^+ \bar{\Lambda}_c^-$ production rates \Rightarrow study $\Lambda_c^+ \rightarrow \pi^+ / K^+ + \nu_s$
- Signature: $e^+e^- \rightarrow \Lambda_c^+ \bar{\Lambda}_c^-$, $\bar{\Lambda}_c^-$ tagging and $\Lambda_c^+ \rightarrow M^+ + \text{missing}$
- Apparent BNV in Λ_c^+ decays at STCF

Model frameworks – the ν LEFT


$$\mathcal{O}_{cdd}^{S,RR} = \epsilon^{\alpha\beta\gamma} (\overline{c}_{R\alpha}^c d_{R\beta}) (\overline{\nu}_L d_{R\gamma})$$

$$\mathcal{O}_{c ds}^{S,RR} = \epsilon^{\alpha\beta\gamma} (\overline{c}_{R\alpha}^c d_{R\beta}) (\overline{\nu}_L s_{R\gamma})$$

$$\mathcal{O}_{csd}^{S,RR} = \epsilon^{\alpha\beta\gamma} (\overline{c}_{R\alpha}^c s_{R\beta}) (\overline{\nu}_L d_{R\gamma})$$

$$\mathcal{L}_{\text{eff.}}^{cdd} \supset \frac{C_{211}}{\Lambda^2} \mathcal{O}_{cdd}^{S,RR} + \text{h.c.}$$

$$\mathcal{L}_{\text{eff.}}^{c ds, csd} \supset \frac{C_{212}}{\Lambda^2} \mathcal{O}_{c ds}^{S,RR} + \frac{C_{221}}{\Lambda^2} \mathcal{O}_{csd}^{S,RR} + \text{h.c.}$$




ν LEFT matched to BChPT

- To compute $\Gamma(\Lambda_c^+ \rightarrow \pi^+/K^+ + \nu_s)$, we follow the literature for **matching the ν LEFT to the BChPT**
- However, BChPT is rigorously applicable only to the **light-flavor baryon octet**, and its direct use for the charmed baryon Λ_c^+ **lies outside its formal domain of validity**
- To account for this mismatch, we **promote** the light-baryon-level (**proton**) matrix elements to the Λ_c^+ case and apply a **conservative order-unity variation** to the corresponding hadronic form factors
- \Rightarrow We vary each Λ_c^+ matrix element by a multiplicative factor of **2 and 1/2** as a **systematic uncertainty** associated with extrapolating the low-energy EFT matching into the heavy-flavor sector
- Captures the expected size of SU(3)-breaking and heavy-quark-symmetry-breaking effects in the absence of a controlled chiral expansion for charm baryons.

The $|\Delta(B-L)| = 2$ effective Lagrangian

$$\mathcal{L}_{\text{eff}} = g_{MB}^{\Lambda_c^+} \bar{B} \gamma^\mu \gamma_5 \Lambda_c^+ \partial_\mu M + m_{Bs,L} \bar{\nu}_s P_R B + i y_{Ms,L}^{\Lambda_c^+} \bar{\nu}_s P_R \Lambda_c^+ M$$



$$\begin{aligned} \overline{|\mathcal{M}|^2}/m_{\Lambda_c^+}^2 &= \frac{1}{2} |y_{Ms,L}^{\Lambda_c^+}|^2 (1 + x_s^2 - x_M^2) + \frac{1}{2} \sum_{B,B'} g_{MB}^{\Lambda_c^+} g_{MB'}^{\Lambda_c^+*} m_{Bs,L} m_{B's,L}^* g(x_B, x_{B'}) \\ &\quad - \sum_B \text{Re}(y_{Ms,L}^{\Lambda_c^+} m_{Bs,L}^* g_{MB}^{\Lambda_c^+*}) h(x_B) \end{aligned}$$

$$x_s = m_{\nu_s}/m_{\Lambda_c^+}, \quad x_M = m_M/m_{\Lambda_c^+}, \quad x_{B^{(\prime)}} = m_{B^{(\prime)}}/m_{\Lambda_c^+}$$

$$\Gamma(\Lambda_c^+ \rightarrow M^+ + \nu_s) = m_{\Lambda_c^+} \frac{\lambda^{1/2}(1, x_M^2, x_s^2)}{16\pi} \frac{\overline{|\mathcal{M}|^2}}{m_{\Lambda_c^+}^2},$$

$$M^+ = \pi^+, K^+$$

The RPV-SUSY




- A light neutralino $\tilde{\chi}_1^0$ being dominantly bino-like

$$\mathcal{L}_{\text{BNV-bino}} = \mathcal{L}_{\text{bino}} + \mathcal{L}_{\text{RPV}}$$

$$\mathcal{L}_{\text{bino}} = - \sum_{q=d,s,c} g_{1R}^{\tilde{q}} (\bar{q}_{R,a} P_L \tilde{\chi}_1^0) \tilde{q}_{R,a} + \text{h.c.} + \dots$$

$$\mathcal{L}_{\text{RPV}} = \lambda_{212}'' \epsilon_{abc} \left(\tilde{c}_{Ra}^* \bar{d}_{Rb} s_{Rc}^C + \tilde{d}_{Ra}^* \bar{c}_{Rb} s_{Rc}^C + \tilde{s}_{Ra}^* \bar{c}_{Rb} d_{Rc}^C \right) + \text{h.c.}$$

$$g_{1R}^{\tilde{q}} = -\sqrt{2} g_W e_q \tan \theta_W$$



The RPV-SUSY matched to the ν LEFT

$$\mathcal{L}^{\text{BNV}} = \frac{\tilde{g}_{1R}^{\tilde{s}} \lambda_{212}''}{m_{\tilde{s}}^2} (\overline{d_R} c_R^c) (\overline{s_R} \tilde{\chi}_1^0) + \frac{\tilde{g}_{1R}^{\tilde{d}} \lambda_{212}''}{m_{\tilde{d}}^2} (\overline{s_R} c_R^c) (\overline{d_R} \tilde{\chi}_1^0) + \frac{\tilde{g}_{1R}^{\tilde{c}} \lambda_{212}''}{m_{\tilde{c}}^2} (\overline{s_R} d_R^c) (\overline{c_R} \tilde{\chi}_1^0)$$

- The first two terms match the h.c. counterparts of $\mathcal{O}_{c\tilde{d}s}^{S,RR}$ and $\mathcal{O}_{c\tilde{s}d}^{S,RR}$
- The third term can be Fierz-transformed:

$$(\overline{s_R} d_R^c) (\overline{c_R} \tilde{\chi}_1^0) \stackrel{\text{Fierz}}{\approx} \frac{1}{2} (\overline{s_R} \tilde{\chi}_1^0) (\overline{c_R} d_R^c) = -\frac{1}{2} (\overline{d_R} c_R^c) (\overline{s_R} \tilde{\chi}_1^0)$$

where we have ignored an extra tensor term that originates from the Fierz transformation and is expected to give negligible contributions

- It matches the h.c. counterpart of $\mathcal{O}_{c\tilde{d}s}^{S,RR}$

$$\Rightarrow \mathcal{L}^{\text{BNV}} = 2 \frac{\tilde{g}_{1R}^{\tilde{s}} \lambda_{212}''}{m_{\tilde{q}}^2} (\overline{d_R} c_R^c) (\overline{s_R} \tilde{\chi}_1^0) + \frac{\tilde{g}_{1R}^{\tilde{d}} \lambda_{212}''}{m_{\tilde{q}}^2} (\overline{s_R} c_R^c) (\overline{d_R} \tilde{\chi}_1^0)$$

- We have assumed squark-mass degeneracy and used $\tilde{g}_{1R}^{\tilde{c}} = -2\tilde{g}_{1R}^{\tilde{s}}$

$$\Rightarrow \frac{c_{212}}{\Lambda^2} = \frac{2\lambda_{212}'' \tilde{g}_{1R}^{\tilde{s}}}{m_{\tilde{q}}^2}, \quad \frac{c_{221}}{\Lambda^2} = \frac{\lambda_{212}'' \tilde{g}_{1R}^{\tilde{d}}}{m_{\tilde{q}}^2}$$

The very long decay lengths of ν_s and $\tilde{\chi}_1^0$

- For the considered mass range of interest, their decays are highly suppressed
- ν_s : decay amplitudes suppressed by the small values of the Wilson coefficients, off-shell propagators of a W and a down-type quark
- $\tilde{\chi}_1^0$: decay amplitudes suppressed by three off-shell propagators (a squark, a W -boson, and a down-type quark), CKM matrix elements, tiny RPV couplings, as well as the absence of squark mixing
- \Rightarrow Both ν_s and $\tilde{\chi}_1^0$ appear as missing energy in the main detector



The STCF and search analysis

- **STCF**: an **energy-symmetry** e^-e^+ collider,
 $E_{\text{COM}} = 4.630 \text{ GeV} \Rightarrow N_{\Lambda_c^+ \bar{\Lambda}_c^-} = 5.6 \times 10^8$ per year with 1 ab^{-1}
- Simulate both signal and background events with **OSCAR**
- $e^+e^- \rightarrow \Lambda_c^+ \bar{\Lambda}_c^-$: **tag** $\bar{\Lambda}_c^-$ and $\Lambda_c^+ \rightarrow M^+ + \nu_s / \tilde{\chi}_1^0$
- **Signature**: $\bar{\Lambda}_c^-$ and $M^+ + \text{missing}$
- What we **have done**:
 - Simulated the tag channel $\bar{\Lambda}_c^- \rightarrow \bar{p}K^+\pi^-$ with $\text{BR} = 6.35\%$, found a reconstruction efficiency of $\sim 39\%$ for the $\Lambda_c^+ \rightarrow M^+$ transition
- **To-do**:
 - Five more tag channels to do: $\bar{\Lambda}\pi^-$, $\bar{\Lambda}\pi^-\pi^0$, $\bar{p}K_S^0$, $\bar{\Sigma}^0\pi^-$, $\bar{\Sigma}^-\pi^-\pi^+$
 - Inclusive background simulations (expected to be low)
 - Selection cuts
- **For now**: use a flat efficiency ϵ of **39%** for all analyses, assume **zero background**, a factor **2.5** to include all tag channels:

$$N_S = 2 \cdot N_{\Lambda_c^+ \bar{\Lambda}_c^-} \cdot \text{BR}(\Lambda_c^+ \rightarrow M^+ + \nu_s / \tilde{\chi}_1^0) \cdot \text{BR}(\bar{\Lambda}_c^- \rightarrow \bar{p}K^+\pi^-) \cdot 2.5 \cdot \epsilon$$


Numerical results – EFT

- Bound on $\text{BR}(\Lambda_c^+ \rightarrow M^+ + \nu_s / \tilde{\chi}_1^0) \sim \mathcal{O}(10^{-8})$



Probe Λ up to 5 – 10 TeV for $c = 1$

Numerical results – RPV-SUSY



Probe λ''_{212} down to $\mathcal{O}(0.1)$ for $m_{\tilde{q}} = 2 \text{ TeV}$

Summary

- Propose to search for a **BNV** signature: $\Lambda_c^+ \rightarrow \pi^+/K^+ + \text{missing}$, predicted in a wide range of BSM scenarios
- $\mathcal{O}(10^8)\Lambda_c^+\bar{\Lambda}_c^-$ events at STCF **per year**
- MC simulations with **OSCAR** to determine acceptance and reconstruction efficiencies of both background and signal events
- **Partly done**: one tag channel for the signal events
- **Promising results** with current estimates: probing Λ up to **5 – 10 TeV**
- **Work to do**: five more tag channels, inclusive backgrounds, and selection cuts

Thank You! 感谢垂听!

Back-up slides

Matching hadronic form factors

| Process | $g_{MB}^{\Lambda_c^+}$ | $m_{Bs,L}$ | $y_{Ms,L}^{\Lambda_c^+}$ |
|---------------------------------------|------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------|
| $\Lambda_c^+ \rightarrow \pi^+ \nu_s$ | $g_{\pi\Sigma_c^0}^{\Lambda_c^+} = \frac{D+F}{f_\pi}$ | $m_{\Sigma_{cs,L}^0} = \frac{c_{211}}{\Lambda^2}(\beta)$ | $y_{\pi s,L}^{\Lambda_c^+} = \frac{c_{211}}{\Lambda^2}(-\frac{\beta}{f_\pi})$ |
| $\Lambda_c^+ \rightarrow K^+ \nu_s$ | $g_{K\Xi_c'^0}^{\Lambda_c^+} = \frac{D-F}{\sqrt{2}f_\pi}$ $g_{K\Xi_c^0}^{\Lambda_c^+} = \frac{D+3F}{\sqrt{6}f_\pi}$ | $m_{\Xi_{cs,L}'^0} = \frac{c_{221}}{\Lambda^2}(\frac{\beta}{\sqrt{2}})$ $m_{\Xi_{cs,L}^0} = \frac{c_{221}}{\Lambda^2}(-\frac{\beta}{\sqrt{6}})$ $+ \frac{c_{212}}{\Lambda^2}(-\beta\sqrt{\frac{2}{3}})$ | $y_{Ks,L}^{\Lambda_c^+} = \frac{c_{212}}{\Lambda^2}(-\frac{\beta}{f_\pi})$ |

Matrix element for $\Lambda_c^+ \rightarrow M^+ + \nu_s$ & Kinematic functions

$$i\mathcal{M} = \overline{u}_{\nu_s} P_R \left(-y_{Ms,L}^{\Lambda_c^+} + \sum_B m_{Bs,L} \frac{\not{k} + m_B}{k^2 - m_B^2} g_{MB}^{\Lambda_c^+} \not{p}_M \gamma_5 \right) u_{\Lambda_c^+}$$

$$g(x_1, x_2) = \frac{(x_s^2 + x_1 x_2)(1 - x_M^2 - x_s^2(2 + x_M^2) + x_s^4) - 2x_s^2 x_M^2(x_1 + x_2)}{(x_1^2 - x_s^2)(x_2^2 - x_s^2)}$$

$$h(x_B) = \frac{x_B(1 - x_M^2 - x_s^2) - x_s^2(1 + x_M^2 - x_s^2)}{x_B^2 - x_s^2}$$

Numerical values of hadronic form factors

$$\beta_{\Lambda_c^+} = 0.835 \times 10^{-2} \text{ GeV}^3$$

deduced in [[Dib, Helo, Lyubovitskij, Neill, Soffer, ZSW 2023](#)] from predictions of QCD sum-rule approaches

$$D = 0.730, F = 0.447, f_\pi = 0.13041 \text{ GeV}$$

Derivation of the eff. Lagrangian from the RPV-SUSY – I

$$\mathcal{L}^{\text{BNV}} = \mathcal{L}^{\text{scd}}\tilde{\chi}_1^0 + \mathcal{L}^{\text{cds}}\tilde{\chi}_1^0 + \mathcal{L}^{\text{dcs}}\tilde{\chi}_1^0$$

$$\mathcal{L}^{q_1 q_2 q_3 \tilde{\chi}_1^0} = \mathcal{O}^{q_1 q_2 q_3} \tilde{\chi}_1^0 + \text{h.c.}$$

$$\mathcal{O}^{q_1 q_2 q_3} = g^{\tilde{q}_1 R} \mathcal{O}_{q_1 q_2 q_3}^{LL}$$

$$g^{\tilde{q} R} = \frac{g_{1R}^{\tilde{q}} \lambda_{212}''}{m_{\tilde{q}}^2}$$

$$\mathcal{O}_{q_1 q_2 q_3}^{LL} = \varepsilon_{abc} (\bar{q}_{3,c} P_L C \bar{q}_{2,b}^T) \bar{q}_{1,a} P_L$$

$C = i\gamma^0\gamma^2$: the charge conjugation matrix.

$$\text{and} \quad \begin{aligned} \mathcal{O}_{cds}^{S,RR} &= (\bar{c}_R^c d_R)(\bar{\nu}_L s_R) \quad \& \text{h.c.} \quad \tilde{\mathcal{O}}_{cds}^{S,RR} = (\bar{d}_R c_R^c)(\bar{s}_R \nu_L) \\ \mathcal{O}_{csd}^{S,RR} &= (\bar{c}_R^c s_R)(\bar{\nu}_L d_R) \quad \tilde{\mathcal{O}}_{csd}^{S,RR} = (\bar{s}_R c_R^c)(\bar{\nu}_L d_R) \end{aligned}$$

Derivation of the eff. Lagrangian from the RPV-SUSY – II

The corresponding effective Lagrangian as

$$\mathcal{L}^{\text{BNV}} = \frac{g_{1R}^{\tilde{s}} \lambda_{212}''}{m_{\tilde{s}}^2} (\overline{d_R} c_R^c) (\overline{s_R} \tilde{\chi}_1^0) + \frac{g_{1R}^{\tilde{d}} \lambda_{212}''}{m_{\tilde{d}}^2} (\overline{s_R} c_R^c) (\overline{d_R} \tilde{\chi}_1^0) + \frac{g_{1R}^{\tilde{c}} \lambda_{212}''}{m_{\tilde{c}}^2} (\overline{s_R} d_R^c) (\overline{c_R} \tilde{\chi}_1^0)$$

The first two terms match with $\tilde{\mathcal{O}}_{cds}^{S,RR}$ and $\tilde{\mathcal{O}}_{csd}^{S,RR}$
 The operator in the third term – Fierz transf.:

$$(\overline{s_R} d_R^c) (\overline{c_R} \tilde{\chi}_1^0) \stackrel{\text{Fierz}}{\approx} \frac{1}{2} (\overline{s_R} \tilde{\chi}_1^0) (\overline{c_R} d_R^c) = -\frac{1}{2} (\overline{d_R} c_R^c) (\overline{s_R} \tilde{\chi}_1^0)$$

where an extra tensor term that originates from the Fierz transformation and is expected to give negligible contributions only, has been ignored

The last expression matches $\tilde{\mathcal{O}}_{cds}^{S,RR}$

$$\mathcal{L}^{\text{BNV}} = 2 \frac{g_{1R}^{\tilde{s}} \lambda_{212}''}{m_{\tilde{q}}^2} (\overline{d_R} c_R^c) (\overline{s_R} \tilde{\chi}_1^0) + \frac{g_{1R}^{\tilde{d}} \lambda_{212}''}{m_{\tilde{q}}^2} (\overline{s_R} c_R^c) (\overline{d_R} \tilde{\chi}_1^0)$$

with squark-mass degeneracy and the relation $g_{1R}^{\tilde{c}} = -2g_{1R}^{\tilde{s}}$

$$\frac{c_{212}}{\Lambda^2} = \frac{2\lambda_{212}'' g_{1R}^{\tilde{s}}}{m_{\tilde{q}}^2}, \quad \frac{c_{221}}{\Lambda^2} = \frac{\lambda_{212}'' g_{1R}^{\tilde{d}}}{m_{\tilde{q}}^2}$$

Preliminary efficiencies

| Miss质量 (GeV) | K^+ , N效率 | Miss质量 (GeV) | π^+ , N效率 |
|-----------------|-------------|-----------------|---------------|
| 0.93827 | 38.82% | 0.93827 | 39.18% |
| 1.110 | 38.98% | 1.11 | 39.62% |
| 1.280 | 39.00% | 1.28 | 40.13% |
| 1.450 | 38.89% | 1.45 | 40.78% |
| 1.620 | 36.95% | 1.62 | 40.26% |
| | | 1.79278 | 39.13% |
| | | 1.935 | 38.38% |
| | | 2.075 | 38.44% |

