





--- A spectral reformulation
of the BBGKY hierarchy

Tsinghua University

arXiv: [2507.14243](#), [2509.23978](#), Xingjian Lu, Shuzhe Shi

Background – Time-space Evolution




Hadronic Gas: Boltzmann Eq.
UrQMD, SMASH

Chemical freezeout out

QGP: A Fluid with Very Low Shear Viscosity
to Entropy Density Ratio
*VISHNU (Ohio State U), MUSIC (McGill U),
CLVisc (CCNU), BESHhydro (Ohio State U),
Trajectum(Utrecht U), vHLE*


Pre-equilibrium

Motivation - Early hydrodynamization Puzzle




Motivation - Early hydrodynamization Puzzle

- **Linear Boltzmann Enough? *No!***
 - **Assume** close to equilibrium
 - **But** QGP is created **far-from-equilibrium**
 - Need consider **Nonlinear effect**



Motivation - Early hydrodynamization Puzzle



- **Linear Boltzmann Enough? No!**
 - **Assume** close to equilibrium
 - **But** QGP is created **far-from-equilibrium**
 - Need consider **Nonlinear effect**
- **Nonlinear Boltzmann Enough? No!**
 - **Assume** dilute and no correlation
 - Break down in pre-equilibrium stage
 - **dense:** $l_{\text{interaction}} \approx l_{\text{mean free length}} \ll L$
 - **with correlation:** $f_2(\phi_a, \phi_b) = f_1(\phi_a)f_1(\phi_b)$
 - Need consider **Correlation**

Pre-equilibrium evolution of QGP

Evolve particle r_i, p_i

No test particle method: QMD

Applicability

$$l_{\text{interaction}} \ll l_{\text{mean free length}} \ll L$$

Event-by-event fluctuations

Many-particle correlations

Test particle method involved: BUU

Applicability

$$l_{\text{interaction}} \approx l_{\text{mean free length}} \ll L$$

(High density)

Evolve the distribution function

Lattice Boltz. for linearized Boltz.

Applicability

$$l_{\text{interaction}} \ll l_{\text{mean free length}} \ll L$$

Near-equilibrium

Accurate evolution of $T^{\mu\nu}$

What if ?

$$l_{\text{interaction}} \approx l_{\text{mean free length}} \ll L$$

far-from-equilibrium

Correlation involve

Map: What we do in the building of nonequilibrium statistical mechanics?

Hamilton Equation/
Schrödinger Equation


Hamilton Equation

$$\frac{dq}{dt} = \frac{\partial \mathcal{H}}{\partial p}, \quad \frac{dp}{dt} = -\frac{\partial \mathcal{H}}{\partial q}$$

Schrödinger equation

$$\hat{H}\Psi = i\hbar \frac{\partial}{\partial t} \Psi$$

Map: What we do in the building of nonequilibrium statistical mechanics?



Liouville's theorem


The distribution function is **constant** along any trajectory **in phase space**.

Liouville equation

$$\frac{\partial P_{(N)}}{\partial t} + \sum_{i=1}^N \left(\frac{\partial P_{(N)}}{\partial q_i} \dot{q}_i + \frac{\partial P_{(N)}}{\partial p_i} \dot{p}_i \right) = 0$$

Full distribution function

Map: What we do in the building of nonequilibrium statistical mechanics?



Liouville equation

$$d_t P_{(N)} = C[\int, P_{(N)}]$$

Full distribution function


BBGKY hierarchy

(Bogoliubov–Born–Green–Kirkwood–Yvon)

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$

Reduced distribution function

Map: What we do in the building of nonequilibrium statistical mechanics?



BBGKY hierarchy

(Bogoliubov–Born–Green–Kirkwood–Yvon)

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$


Reduced distribution function

Assume $P_2(\phi_a, \phi_b) - P_1(\phi_a)P_1(\phi_b) = 0$

Boltzmann Equation

$$d_t P_{(1)} = C[\int, P_{(1)}]$$

Map: What we do in the building of nonequilibrium statistical mechanics?



Boltzmann Equation

$$d_t f_1 \propto (f_3 f_4 - f_1 f_2)$$

Around equilibrium

Linear Boltzmann Equation

$$d_t f_1 \propto (f_3 + f_4 - f_1 - f_2)$$


RTA Boltzmann Equation

$$d_t f_1 \propto (f_1 - f_{eq})/\tau$$

Hydrodynamics

$$\nabla_\mu T^{\mu\nu} = 0$$

Map: What we do in the building of nonequilibrium statistical mechanics?



**Time-reversal symmetry,
Cornerstone**



Gap: correlation

Tractable proxy




Gap: nonlinear

Workhorse



Map: What we do in the building of nonequilibrium statistical mechanics?



New theoretical tool: Spectral BBGKY Hierarchy

BBGKY Hierarchy

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$



$\int$ means high dimensional-integral

New theoretical tool: Spectral BBGKY Hierarchy

Difficult to solve

if Evolve the distribution function itself

- **Until now**, almost all simulations are based on **lowest-order truncation**, *the Boltzmann equation*.

BBGKY Hierarchy

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$



$\int$ means high dimensional-integral

New theoretical tool: Spectral BBGKY Hierarchy

Difficult to solve

if Evolve the distribution function itself

- **Until now**, almost all simulations are based on **lowest-order truncation**, the Boltzmann equation.
- Because **huge memory demands** N^{6n} to discretize $6n$ D
 - $n = 1$, discretize $6 D$
 - $n = 2$, discretize $12 D$
 - for ref, Lattice QCD discretize $3 + 1 D$, ~30 GB memory

BBGKY Hierarchy

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$



$\int$ means high dimensional-integral

New theoretical tool: Spectral BBGKY Hierarchy

Spectrum Method

$$P_{(n)}(\mathbf{p}_1, \dots, \mathbf{p}_n) = \underbrace{\mathbf{p}^{i_1 \dots i_n}}_{\text{coefficient}} \underbrace{\mathcal{P}_{i_1}(p_1) \dots \mathcal{P}_{i_n}(p_n)}_{\text{basis}}$$

coefficient

basis

BBGKY Hierarchy

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$

New theoretical tool: Spectral BBGKY Hierarchy

Spectrum Method

$$P_{(n)}(\mathbf{p}_1, \dots, \mathbf{p}_n) = \underbrace{P^{i_1 \dots i_n}}_{\text{coefficient}} \underbrace{\mathcal{P}_{i_1}(p_1) \dots \mathcal{P}_{i_n}(p_n)}_{\text{basis}}$$

coefficient

basis

- not a new idea. But we **basis depends on physics problem.**

- We hope

first few basis functions

already capture the important physics

BBGKY Hierarchy

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$

New theoretical tool: Spectral BBGKY Hierarchy

Spectrum Method

$$P_{(n)}(\mathbf{p}_1, \dots, \mathbf{p}_n) = \underbrace{\mathbf{P}^{i_1 \dots i_n}}_{\text{coefficient}} \underbrace{\mathcal{P}_{i_1}(p_1) \dots \mathcal{P}_{i_n}(p_n)}_{\text{basis}}$$

coefficient

basis

Our basis

$$\mathcal{P}_{n,\ell,m}(p_\mu) = e^{-p_\mu u^\mu / \Lambda} \left(\frac{p_\mu u^\mu}{\Lambda} \right)^\ell Y_{\ell,m}(\theta, \phi) L_n^{(2\ell+2)} \left(\frac{p_\mu u^\mu}{\Lambda} \right)$$

The physics

Angular

Radial

- equilibrium
- particle number, energy-momentum conservation
- nonlinear behavior

mainly depends on the first few basis.

BBGKY Hierarchy

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$

New theoretical tool: Spectral BBGKY Hierarchy

Spectrum Method

$$P_{(n)}(\mathbf{p}_1, \dots, \mathbf{p}_n) = \mathbf{P}^{i_1 \dots i_n} \mathcal{P}_{i_1}(p_1) \dots \mathcal{P}_{i_n}(p_n)$$

BBGKY Hierarchy

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$

$$\begin{aligned} n &= \frac{4\sqrt{\pi}}{(2\pi)^3} T^3 f^{(0,0,0)} \\ \langle p^0 \rangle &= \frac{12\sqrt{\pi}}{(2\pi)^3} T^4 \left(f^{(0,0,0)} - f^{(1,0,0)} \right) \\ \langle p^1 \rangle &= \frac{16\sqrt{3\pi}}{(2\pi)^3} T^4 f^{(0,1,1)} \\ \langle p^2 \rangle &= \frac{16\sqrt{3\pi}}{(2\pi)^3} T^4 f^{(0,1,-1)} \\ \langle p^3 \rangle &= \frac{16\sqrt{3\pi}}{(2\pi)^3} T^4 f^{(0,1,0)} \end{aligned}$$

Our basis

$$\mathcal{P}_{n,\ell,m}(p_\mu) = e^{-p_\mu u^\mu / \Lambda} \left(\frac{p_\mu u^\mu}{\Lambda} \right)^\ell Y_{\ell,m}(\theta, \phi) L_n^{(2\ell+2)} \left(\frac{p_\mu u^\mu}{\Lambda} \right)$$

The physics

- equilibrium
- particle number, energy-momentum conservation
- nonlinear behavior

mainly depends on the first few basis.

Angular Radial

New theoretical tool: Spectral BBGKY Hierarchy

Spectrum Method

$$P_{(n)}(\mathbf{p}_1, \dots, \mathbf{p}_n) = \mathbf{P}^{i_1 \dots i_n} \mathcal{P}_{i_1}(p_1) \dots \mathcal{P}_{i_n}(p_n)$$

BBGKY Hierarchy

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$

Spectral BBGKY Hierarchy

$$\underline{d_t \mathbf{P}_{(n)}} = \underline{\mathbf{C}} \cdot \underline{\mathbf{P}_{(n)}} + \underline{\mathbf{A}} \cdot \underline{\mathbf{P}_{(n+1)}}$$

$\mathbf{A}$ & $\mathbf{C}$: Collision Integral tensor

coefficients

Advantage: Step toward high-density pair-correlation function

BBGKY Hierarchy

$$d_t \underline{P_{(n)}} = C[\underline{\int}, \underline{P_{(n)}}, \underline{P_{(n+1)}}]$$

distributions

Spectral BBGKY Hierarchy

$$\underline{d_t P_{(n)}} = \underline{C} \cdot \underline{P_{(n)}} + \underline{A} \cdot \underline{P_{(n+1)}}$$

A & C : Collision Integral

coefficients

Advantage: Step toward high-density pair-correlation function

BBGKY Hierarchy

$$d_t \underline{P_{(n)}} = C[\underline{\int}, \underline{P_{(n)}}, \underline{P_{(n+1)}}]$$

distributions

Spectral BBGKY Hierarchy

$$\underline{d_t P_{(n)}} = \underline{C} \cdot \underline{P_{(n)}} + \underline{A} \cdot \underline{P_{(n+1)}}$$

A & C : Collision Integral

coefficients

- **BBGKY:** discretize $6n$ D, cost N^{6n}
(3coorediate,3momentum)
- **sBBGKY:** discretize $3n$ D, cost N^{3n}
(3coorediate)

Advantage: Step toward high-density pair-correlation function

BBGKY Hierarchy

$$d_t \underline{P_{(n)}} = C[\underline{\int}, \underline{P_{(n)}}, \underline{P_{(n+1)}}]$$

distributions

Spectral BBGKY Hierarchy

$$\underline{d_t P_{(n)}} = \underline{C} \cdot \underline{P_{(n)}} + \underline{A} \cdot \underline{P_{(n+1)}}$$

A & C : Collision Integral

coefficients

- **before:** discretize $6n$ D, cost N^{6n}
(3coorediate,3momentum)
- **now:** discretize $3n$ D, cost N^{3n}
(3coorediate)

$$\textit{Cost}_{\text{now}} = \sqrt{\textit{Cost}_{\text{before}}}$$

Advantage: No numerical fluctuations from Monte Carlo integration

BBGKY Hierarchy

$$d_t P_{(n)} = C[\underline{\int}, P_{(n)}, P_{(n+1)}]$$

Collision integral coupled time evolution

- Monte Carlo integration
- Numerical fluctuations,
need ensemble average

Spectral BBGKY Hierarchy

$$d_t \mathbf{P}_{(n)} = \mathbf{C} \cdot \mathbf{P}_{(n)} + \mathbf{A} \cdot \mathbf{P}_{(n+1)}$$

$\mathbf{A}$ & $\mathbf{C}$: Collision Integral

Advantage: No numerical fluctuations from Monte Carlo integration

BBGKY Hierarchy

$$d_t P_{(n)} = C[\int, P_{(n)}, P_{(n+1)}]$$

Collision integral coupled time evolution

Spectral BBGKY Hierarchy

$$d_t \mathbf{P}_{(n)} = \underline{\mathbf{C}} \cdot \mathbf{P}_{(n)} + \underline{\mathbf{A}} \cdot \mathbf{P}_{(n+1)}$$

$\mathbf{A}$ & $\mathbf{C}$: Collision Integral

Decouple collision integral from evolution

$$C_{ijks}^{\text{gain}} = \int_{p_{1\mu}, p_{2\mu}} \int_{\mathbf{p}_3, \mathbf{p}_4} Q_i(p_{1\mu}) Q_j(p_{2\mu}) \frac{1}{p_{1\mu}^0 p_{2\mu}^0} \mathcal{W}_{(\mathbf{p}_1, \mathbf{p}_2 \rightarrow \mathbf{p}_3, \mathbf{p}_4)} \mathcal{P}_j(p'_{1\mu}) \mathcal{P}_k(p'_{2\mu})$$

➤ Simplify

8D
Integral

3D
Integral

$m=0$ case
polynomial σ

Finite Sums

➤ Compact


Parity + rotation symmetry

- Analytically calculate Collision Integral
- One simulation per initial state

$$\textit{Cost}_{nonlin} = \textit{Cost}_{lin}$$

No numerical fluctuations

Numerical validation



Hydrodynamization from the full Boltzmann Eq.

We need:

hydrodynamization
= **near equilibrium**

People before

hydrodynamization
= **equilibrium**

Now we can determine

near equilibrium
through linearization

Hydrodynamization from the full Boltzmann Eq.

We need:

hydrodynamization
= **near equilibrium**

People before


hydrodynamization
= **equilibrium**

Now we can determine

near equilibrium
through linearization

$$\mathcal{M}_{\Pi}(t) \equiv \frac{4\pi^{\frac{5}{2}}}{T^2} \int \frac{d^3\mathbf{p}}{(2\pi)^3 p^0} (f(t, \mathbf{p}) - f_{\text{eq}}(t, \mathbf{p}))$$

Hydrodynamization from the full Boltzmann Eq.



$$\mathcal{M}_{\Pi}(t) \equiv \frac{4\pi^{\frac{5}{2}}}{T^2} \int \frac{d^3\mathbf{p}}{(2\pi)^3 p^0} (f(t, \mathbf{p}) - f_{\text{eq}}(t, \mathbf{p}))$$

We need:

hydrodynamization
= **near equilibrium**


People before

hydrodynamization
= **equilibrium**

Now we can determine

near equilibrium
through linearization

Hydrodynamization from the full Boltzmann Eq.



$$\mathcal{M}_{\Pi}(t) \equiv \frac{4\pi^{\frac{5}{2}}}{T^2} \int \frac{d^3\mathbf{p}}{(2\pi)^3 p^0} (f(t, \mathbf{p}) - f_{\text{eq}}(t, \mathbf{p}))$$

We need:

hydrodynamization
= **near equilibrium**

People before

hydrodynamization
= **equilibrium**


Now we can determine

near equilibrium
through linearization

- Under isotropic conditions, a **nonlinear** system already enters a **linear** regime before it approaches **thermal equilibrium**

$$\tau_{\text{lin}}^{[O]} < \tau_{\text{therm}}^{[O]}$$

Hydrodynamization from the full Boltzmann Eq.



We need:

hydrodynamization
= **near equilibrium**


People before

hydrodynamization
= **equilibrium**

Now we can determine

near equilibrium
through linearization

Hydrodynamization from the full Boltzmann Eq.



We need:

hydrodynamization
= **near equilibrium**

People before


hydrodynamization
= **equilibrium**

Now we can determine

near equilibrium
through linearization

“distribution ensemble”

Hydrodynamization from the full Boltzmann Eq.



We need:

hydrodynamization
= **near equilibrium**


People before

hydrodynamization
= **equilibrium**

Now we can determine

near equilibrium
through linearization

Hydrodynamization from the full Boltzmann Eq.



We need:

hydrodynamization
= **near equilibrium**

People before

hydrodynamization
= **equilibrium**

Now we can determine

near equilibrium
through linearization


- Observables

$$\mathcal{M}_{\Pi} = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{E} f(t, \mathbf{p})$$

$$\mathcal{M}_i = \int \frac{d^3 p}{(2\pi)^3} E^i f(t, \mathbf{p}) \quad i = 0, 1, \dots, 10$$

look at the blue bands

Hydrodynamization from the full Boltzmann Eq.



We need:

hydrodynamization
= **near equilibrium**

People before

hydrodynamization
= **equilibrium**

Now we can determine

near equilibrium
through linearization

• Observables


$$\mathcal{M}_{\Pi} = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{E} f(t, \mathbf{p})$$

$$\mathcal{M}_i = \int \frac{d^3 p}{(2\pi)^3} E^i f(t, \mathbf{p}) \quad i = 0, 1, \dots, 10$$

- Different observables share **similar** thermalization and linearization times

$$\tau_{\text{lin}}^{[\sigma(\mathcal{M}_{\Pi})]} \approx \tau_{\text{lin}}^{[\sigma(\mathcal{M}_s)]}, \quad \tau_{\text{therm}}^{[\sigma(\mathcal{M}_{\Pi})]} \approx \tau_{\text{therm}}^{[\sigma(\mathcal{M}_s)]}$$

Hydrodynamization from the full Boltzmann Equation




➔ Both thermalization and linearization have slowest

theoretical limits,


$$\text{upper limit } \tau_{\text{lin}}^{[\sigma(\mathcal{M}_s)]} = \frac{1}{2} \text{ upper limit } \tau_{\text{therm}}^{[\sigma(\mathcal{M}_s)]}$$

look at the orange lines

Hydrodynamization from the full Boltzmann Equation



Hydrodynamization from the full Boltzmann Equation




Independent of the initial distribution ensembles

$$\frac{\delta\tau^{[\sigma(\mathcal{M})]}_{\text{therm}}}{\delta f_{\text{init}}} = 0, \quad \frac{\delta\tau^{[\sigma(\mathcal{M})]}_{\text{lin}}}{\delta f_{\text{init}}} = 0$$

Hydrodynamization from the full Boltzmann equation

Moment $\mathcal{M}_{i,j,k,s}(t) = \int d^3\mathbf{p} f(t, \mathbf{p}) (E_{\mathbf{p}})^i (p^x)^j (p^y)^k (p^z)^s$

$T^{\mu\nu} \quad T^{\mu\nu} = \int_{\mathbf{p}} p^\mu p^\nu f(t, \mathbf{x}, \mathbf{p})$




Consistent: $\tau_{\text{lin}} \approx 0.5 \tau_{\text{therm}}$

Hydrodynamization from the full Boltzmann equation

Moment

$$\mathcal{M}_{i,j,k,s}(t) = \int d^3\mathbf{p} f(t, \mathbf{p}) (E_{\mathbf{p}})^i (p^x)^j (p^y)^k (p^z)^s$$

$$\mathbf{T}^{\mu\nu} \quad T^{\mu\nu} = \int_{\mathbf{p}} p^\mu p^\nu f(t, \mathbf{x}, \mathbf{p})$$



Consistent: $\tau_{\text{lin}} \approx 0.5 \tau_{\text{therm}}$


Summary

- New theoretical tool: **Spectral BBGKY Hierarchy**(arXiv:2509.23978)
 - **Fill the Gap: Correlation, Nonlinear**
- Physical Insight: Decoupling hydrodynamization from thermalization via **nonlinear** Boltzmann equation(arXiv:2509.23978)

$$\tau_{lin} \approx \tau_{therm}/2$$

Outlook

- The thermalization time and linearization time in an expanding system
- The correlation contribution to thermalization process
- $gg \leftrightarrow ggg$
- Bose Enhancement, Pauli Blocking



From Framework to Computation: How We Actually Solve It

momentum expansion

evaluation of the collision integral

BBGKY

Bogoliubov, Born, Green, Kirkwood, Yvon

Liouville Equation

$$\frac{\partial P_{(N)}(\{\phi\})}{\partial t} + \sum_{i=1}^N \frac{\mathbf{p}_i}{p_i^0} \cdot \frac{\partial P_{(N)}(\{\phi\})}{\partial \mathbf{x}_i} = \int d^{6N} \{\phi'\} \Gamma_{\{\phi\} \rightarrow \{\phi'\}} \left(P_{(N)}(\{\phi'\}) - P_{(N)}(\{\phi\}) \right)$$

Interaction

$$= \sum_{\substack{i,j=1 \\ i < j}}^N \mathcal{W}_{\mathbf{p}_i, \mathbf{p}_j \rightarrow \mathbf{p}'_i, \mathbf{p}'_j} \delta^{(3)}(\mathbf{x}_i - \mathbf{x}_j) \delta^{(3)}(\mathbf{x}'_i - \mathbf{x}'_j) \delta^{(3)}(\mathbf{x}_i - \mathbf{x}'_i) \left(\prod_{k \neq i,j} \delta^{(6)}(\phi_k - \phi'_k) \right)$$

$$\begin{aligned} & \frac{\partial P_{(n)}(\{\phi\})}{\partial t} + \sum_{i=1}^n \frac{\mathbf{p}_i}{p_i^0} \cdot \frac{\partial P_{(n)}(\{\phi\})}{\partial \mathbf{x}_i} \\ &= \sum_{\substack{i,j=1 \\ i < j}}^n \delta^{(3)}(\mathbf{x}_i - \mathbf{x}_j) \frac{1}{p_i^0 p_j^0} \int_{\mathbf{p}'_i, \mathbf{p}'_j} \mathcal{W}_{\mathbf{p}_i, \mathbf{p}_j \rightarrow \mathbf{p}'_i, \mathbf{p}'_j} \left(P_{(n)}(\mathbf{p}'_i, \mathbf{p}'_j, \dots) - P_{(n)}(\mathbf{p}_i, \mathbf{p}_j, \dots) \right) \\ &+ (N - n) \sum_{i=1}^n \frac{1}{p_i^0} \int_{\mathbf{p}'_i, \mathbf{p}'_j, \mathbf{p}_j} \mathcal{W}_{\mathbf{p}_i, \mathbf{p}_j \rightarrow \mathbf{p}'_i, \mathbf{p}'_j} \left(P_{(n+1)}(\mathbf{p}'_i, \dots; \mathbf{p}'_j) - P_{(n+1)}(\mathbf{p}_i, \dots; \mathbf{p}_j) \right) \end{aligned}$$

BBGKY

Spectral BBGKY

Spectral BBGKY

$$\begin{aligned}
 & \frac{\partial}{\partial t} \mathbf{P}^{t_1 t_2 \cdots t_n}(t, \mathbf{x}_1, \mathbf{x}_2, \cdots, \mathbf{x}_n) \\
 & + \sum_{j=1}^n \mathbf{B}_{t_j h_j} \cdot \frac{\partial}{\partial \mathbf{x}_j} \mathbf{P}^{t_1 t_2 \cdots h_j \cdots t_n}(t, \mathbf{x}_1, \mathbf{x}_2, \cdots, \mathbf{x}_n) \\
 & = \sum_{\substack{i,j=1 \\ i < j}}^n \delta^{(3)}(\mathbf{x}_i - \mathbf{x}_j) C_{t_i t_j h_i h_j} \mathbf{P}^{t_1 t_2 \cdots h_i \cdots h_j \cdots t_n}(t, \mathbf{x}_1, \mathbf{x}_2, \cdots, \mathbf{x}_n) \\
 & + (N - n) \sum_{i=1}^n A_{t_i h_i h_j} \mathbf{P}^{t_1 t_2 \cdots h_i \cdots t_n h_j}(t, \mathbf{x}_1, \mathbf{x}_2, \cdots, \mathbf{x}_i, \cdots, \mathbf{x}_n, \mathbf{x}_i)
 \end{aligned}$$

$$\mathbf{B}_{ij} = \int_{p_\mu} \mathcal{Q}_i(p_\mu) \mathcal{P}_j(p_\mu) \frac{\mathbf{p}}{p^0}$$

$$A_{ijk} = A_{ijk}^{\text{gain}} - A_{ijk}^{\text{loss}},$$

$$C_{ijks} = C_{ijks}^{\text{gain}} - C_{ijks}^{\text{loss}},$$

$$\mathcal{P}_{n,\ell,m}(p_\mu) = e^{-p_\mu u^\mu / \Lambda} \left(\frac{p_\mu u^\mu}{\Lambda} \right)^\ell Y_{\ell,m}(\theta, \phi) L_n^{(2\ell+2)} \left(\frac{p_\mu u^\mu}{\Lambda} \right)$$

$$\mathcal{Q}_{n,\ell,m}(p_\mu) = \frac{n!}{\Gamma(2\ell + n + 3)} \left(\frac{p_\mu u^\mu}{\Lambda} \right)^{\ell+2} Y_{\ell,m}(\theta, \phi) L_n^{(2\ell+2)} \left(\frac{p_\mu u^\mu}{\Lambda} \right)$$

$$A_{ijk}^{\text{gain}} = \int_{p_{1\mu}} \int_{\mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4} \mathcal{Q}_i(p_{1\mu}) \frac{1}{p_1^0} \mathcal{W}_{(\mathbf{p}_1, \mathbf{p}_2 \rightarrow \mathbf{p}_3, \mathbf{p}_4)} \mathcal{P}_j(p'_{1\mu}) \mathcal{P}_k(p'_{2\mu}),$$

$$A_{ijk}^{\text{loss}} = \int_{p_{1\mu}} \int_{\mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4} \mathcal{Q}_i(p_{1\mu}) \frac{1}{p_1^0} \mathcal{W}_{(\mathbf{p}_1, \mathbf{p}_2 \rightarrow \mathbf{p}_3, \mathbf{p}_4)} \mathcal{P}_j(p_{1\mu}) \mathcal{P}_k(p_{2\mu}),$$

$$C_{ijks}^{\text{gain}} = \int_{p_{1\mu}, p_{2\mu}} \int_{\mathbf{p}_3, \mathbf{p}_4} \mathcal{Q}_i(p_{1\mu}) \mathcal{Q}_j(p_{2\mu}) \frac{1}{p_1^0 p_2^0} \mathcal{W}_{(\mathbf{p}_1, \mathbf{p}_2 \rightarrow \mathbf{p}_3, \mathbf{p}_4)} \mathcal{P}_j(p'_{1\mu}) \mathcal{P}_k(p'_{2\mu})$$

$$C_{ijks}^{\text{loss}} = \int_{p_{1\mu}, p_{2\mu}} \int_{\mathbf{p}_3, \mathbf{p}_4} \mathcal{Q}_i(p_{1\mu}) \mathcal{Q}_j(p_{2\mu}) \frac{1}{p_1^0 p_2^0} \mathcal{W}_{(\mathbf{p}_1, \mathbf{p}_2 \rightarrow \mathbf{p}_3, \mathbf{p}_4)} \mathcal{P}_j(p_{1\mu}) \mathcal{P}_k(p_{2\mu})$$

Minimizing Collision Integral Evaluations

$$A_{ijk}, C_{ijks}$$

- Reduction relationship

$$C_{i0\langle ks\rangle} = C_{0i\langle ks\rangle} = \frac{(2\pi)^3}{\sqrt{4\pi}} \frac{1}{T^3} A_{i\langle ks\rangle}$$

- Parity symmetry

$$P Y_\ell^m(\theta, \varphi) P = (-1)^\ell Y_\ell^m(\theta, \varphi) \longrightarrow C_{ijks} = 0, \quad \text{if } \ell_i + \ell_j + \ell_k + \ell_s \text{ is odd}$$

- Rotational invariance

$$Y_\ell^m(\theta', \varphi') = \sum_{m'=-\ell}^{\ell} \left[D_{mm'}^{(\ell)}(\mathcal{R}) \right]^* Y_\ell^{m'}(\theta, \varphi) \longrightarrow 0 = (\mathcal{D}_{ijks, i'j'k's'} - I) C_{i'j'k's'}$$

$$Y_\ell^m(\theta, \varphi + \Delta\varphi) = e^{im\Delta\varphi} Y_\ell^m(\theta, \varphi) \longrightarrow C_{ijks} = 0, \quad \text{if } m_i + m_j \neq m_k + m_s$$

$$n \in \{0, 1, 2\}, l \in \{0, 1, 2, 3\} \quad 2 \times 48^4 = 1.06 \times 10^7 \xrightarrow{0.2\%} 26244$$

Integral Form of the Collision Kernel

Gernal Cases

8 Fold



3 Fold

$$\begin{aligned}
 & A_{n\ell m, n_1\ell_1 m_1, n_2\ell_2 m_2, n_3\ell_3 m_3, n_4\ell_4 m_4} \\
 &= \frac{n!}{\Gamma(2\ell + n + 3)} \frac{S_{(\mu_1 \dots \mu_\ell)}^{*(m)} S_{(\alpha_1 \dots \alpha_{\ell_1})}^{(m_1)} S_{(\varepsilon_1 \dots \varepsilon_{\ell_2})}^{(m_2)} S_{(\omega_1 \dots \omega_{\ell_3})}^{(m_3)} S_{(\lambda_1 \dots \lambda_{\ell_4})}^{(m_4)}}{(2\pi)^5 \Lambda^{\ell + \ell_1 + \ell_2 + \ell_3 + \ell_4 + 5}} \\
 &\times \int d|\mathbf{P}| |\mathbf{P}|^2 \int_{|\mathbf{P}|}^\infty dP^0 P^2 e^{-2P^\mu u_\mu / \Lambda} \frac{\sqrt{-Q^2}}{2P} \frac{\sqrt{-Q'^2}}{2P} \sum_{g=0}^\infty \sigma^{(g)} \left(\frac{P}{\Lambda} \right) \frac{4\pi}{2g+1} \sum_{h=-g}^g \\
 &\times \left[\int d\Omega_P \left(\prod_{i=1}^\ell \mathcal{M}_{\nu_i}^{\mu_i} \right) \left(\prod_{i=1}^{\ell_1} \mathcal{M}_{\beta_i}^{\alpha_i} \right) \left(\prod_{i=1}^{\ell_2} \mathcal{M}_{\zeta_i}^{\varepsilon_i} \right) \left(\prod_{i=1}^{\ell_3} \mathcal{M}_{\kappa_i}^{\omega_i} \right) \left(\prod_{i=1}^{\ell_4} \mathcal{M}_{\rho_i}^{\lambda_i} \right) \right] \\
 &\times \int_0^\pi \sin \theta d\theta \left(\frac{p_{1\mu} u^\mu}{|\mathbf{p}_1|} \right)^{\ell + \ell_1 + 1} \left(\frac{p_{2\mu} u^\mu}{|\mathbf{p}_2|} \right)^{\ell_2} L_n^{(2\ell+2)} \left(\frac{p_{1\mu} u^\mu}{\Lambda} \right) L_{n_1}^{(2\ell_1+2)} \left(\frac{p_{1\mu} u^\mu}{\Lambda} \right) L_{n_2}^{(2\ell_2+2)} \left(\frac{p_{2\mu} u^\mu}{\Lambda} \right) \\
 &\times \int_0^\pi \sin \theta' d\theta' \left(\frac{p_{3\mu} u^\mu}{|\mathbf{p}_3|} \right)^{\ell_3} \left(\frac{p_{4\mu} u^\mu}{|\mathbf{p}_4|} \right)^{\ell_4} L_{n_3}^{(2\ell_3+2)} \left(\frac{p_{3\mu} u^\mu}{\Lambda} \right) L_{n_4}^{(2\ell_4+2)} \left(\frac{p_{4\mu} u^\mu}{\Lambda} \right) \\
 &\times \left[\int_0^{2\pi} d\phi \left(\prod_{i=1}^\ell p_1^{\nu_i} \right) \left(\prod_{i=1}^{\ell_1} p_1^{\beta_i} \right) \left(\prod_{i=1}^{\ell_2} p_2^{\zeta_i} \right) Y_g^{*h}(\theta, \phi) \right] \left[\int_0^{2\pi} d\phi' \left(\prod_{i=1}^{\ell_3} p_3^{\kappa_i} \right) \left(\prod_{i=1}^{\ell_4} p_4^{\rho_i} \right) Y_g^h(\theta', \phi') \right].
 \end{aligned}$$

Integral Form of the Collision Kernel

Gernal Cases

8 Fold



3 Fold

$$\begin{aligned}
 & C_{n_a l_a m_a, n_b l_b m_b, n_1 l_1 m_1, n_2 l_2 m_2, n_3 l_3 m_3, n_4 l_4 m_4} \\
 = & \frac{n_a!}{\Gamma(2\ell_a + n_a + 3)} \frac{n_b!}{\Gamma(2\ell_b + n_b + 3)} \frac{S_{(\mu_1 \dots \mu_{\ell_a})}^{*(m_a)} S_{(\eta_1 \dots \eta_{\ell_b})}^{*(m_b)} S_{(\alpha_1 \dots \alpha_{\ell_1})}^{(m_1)} S_{(\varepsilon_1 \dots \varepsilon_{\ell_2})}^{(m_2)} S_{(\omega_1 \dots \omega_{\ell_3})}^{(m_3)} S_{(\lambda_1 \dots \lambda_{\ell_4})}^{(m_4)}}{(2\pi)^2 \Lambda^{\ell_a + \ell_b + \ell_1 + \ell_2 + \ell_3 + \ell_4 + 8}} \\
 & \times \int d|\mathbf{P}| |\mathbf{P}|^2 \int_{|\mathbf{P}|}^{\infty} dP^0 P^2 e^{-2P^\mu u_\mu / \Lambda} \frac{\sqrt{-Q^2}}{2P} \frac{\sqrt{-Q'^2}}{2P} \sum_{g=0}^{\infty} \sigma^{(g)} \left(\frac{P}{\Lambda} \right) \frac{4\pi}{2g+1} \sum_{h=-g}^g \\
 & \times \left[\int d\Omega_P \left(\prod_{k=1}^{\ell_a} \mathcal{M}_{\nu_k}^{\mu_k} \right) \left(\prod_{k=1}^{\ell_b} \mathcal{M}_{\chi_k}^{\eta_k} \right) \left(\prod_{k=1}^{\ell_1} \mathcal{M}_{\beta_k}^{\alpha_k} \right) \left(\prod_{k=1}^{\ell_2} \mathcal{M}_{\zeta_k}^{\varepsilon_k} \right) \left(\prod_{k=1}^{\ell_3} \mathcal{M}_{\kappa_k}^{\omega_k} \right) \left(\prod_{k=1}^{\ell_4} \mathcal{M}_{\rho_k}^{\lambda_k} \right) \right] \\
 & \times \int_0^\pi \sin \theta d\theta \left(\frac{p_{1\mu} u^\mu}{|\mathbf{p}_1|} \right)^{\ell_a + \ell_1 + 1} \left(\frac{p_{2\mu} u^\mu}{|\mathbf{p}_2|} \right)^{\ell_b + \ell_2 + 1} \\
 & \times L_{n_a}^{(2\ell_a + 2)} \left(\frac{p_{1\mu} u^\mu}{\Lambda} \right) L_{n_b}^{(2\ell_b + 2)} \left(\frac{p_{2\mu} u^\mu}{\Lambda} \right) L_{n_1}^{(2\ell_1 + 2)} \left(\frac{p_{1\mu} u^\mu}{\Lambda} \right) L_{n_2}^{(2\ell_2 + 2)} \left(\frac{p_{2\mu} u^\mu}{\Lambda} \right) \\
 & \times \int_0^\pi \sin \theta' d\theta' \left(\frac{p_{3\mu} u^\mu}{|\mathbf{p}_3|} \right)^{\ell_3} \left(\frac{p_{4\mu} u^\mu}{|\mathbf{p}_4|} \right)^{\ell_4} L_{n_3}^{(2\ell_3 + 2)} \left(\frac{p_{3\mu} u^\mu}{\Lambda} \right) L_{n_4}^{(2\ell_4 + 2)} \left(\frac{p_{4\mu} u^\mu}{\Lambda} \right) \\
 & \times \left[\int_0^{2\pi} d\phi \left(\prod_{k=1}^{\ell_i} p_1^{\nu_k} \right) \left(\prod_{k=1}^{\ell_j} p_2^{\chi_k} \right) \left(\prod_{i=1}^{\ell_1} p_1^{\beta_i} \right) \left(\prod_{i=1}^{\ell_2} p_2^{\zeta_i} \right) Y_g^{*h}(\theta, \phi) \right] \\
 & \times \left[\int_0^{2\pi} d\phi' \left(\prod_{i=1}^{\ell_3} p_3^{\kappa_i} \right) \left(\prod_{i=1}^{\ell_4} p_4^{\rho_i} \right) Y_g^h(\theta', \phi') \right].
 \end{aligned}$$

Summation Form of the Collision Kernel

$$\begin{aligned}
 & \frac{A_{n_a \ell_a 0, n_1 \ell_1 0, n_2 \ell_2 0, n_3 \ell_3 0, n_4 \ell_4 0}}{\Gamma(2\ell_a + n_a + 3)} \prod_{i \in \{a, 1, 2, 3, 4\}} \sqrt{2\ell_i + 1} \\
 & \sum_{w'_i=0}^{\lfloor \frac{\ell_i}{2} \rfloor + \lfloor \frac{\ell_i}{2} \rfloor} \sum_{\eta'_i=0}^{w'_i - \eta'_i} \sum_{\epsilon'_i=0}^{w'_i - \eta'_i} \sum_{f'_i=0}^{n_a + n_1} \left[\prod_{i \in \{2, 3, 4\}} \sum_{w_i=0}^{\lfloor \frac{\ell_i}{2} \rfloor} \sum_{\eta_i=0}^{w_i - \eta_i} \sum_{\epsilon_i=0}^{w_i - \eta_i} \sum_{f_i=0}^{n_i} \right] \sum_{t'_{12}=0}^{f'_1 + f_2} \sum_{t_{34}=0}^{f_3 + f_4} \\
 & \sum_{x'_{i,12} + x'_{j,12} + x'_{k,12} \leq 2\eta'_{12}} \sum_{y'_{i,12} + y'_{j,12} + y'_{k,12} \leq 2\epsilon'_{12}} \sum_{z'_{i,12} + z'_{k,12} \leq \ell'_{12} - 2\eta'_{12} - 2\epsilon'_{12}} \\
 & \sum_{x'_{i,12}, x'_{j,12}, x'_{k,12} = 0} \sum_{y'_{i,12}, y'_{j,12}, y'_{k,12} = 0} \sum_{z'_{i,12}, z'_{k,12} = 0} \\
 & \sum_{x_{i,34} + x_{j,34} + x_{k,34} \leq 2\eta_{34}} \sum_{y_{i,34} + y_{j,34} + y_{k,34} \leq 2\epsilon_{34}} \sum_{z_{i,34} + z_{k,34} \leq \ell_{34} - 2\eta_{34} - 2\epsilon_{34}} \\
 & \sum_{x_{i,34}, x_{j,34}, x_{k,34} = 0} \sum_{y_{i,34}, y_{j,34}, y_{k,34} = 0} \sum_{z_{i,34}, z_{k,34} = 0} \\
 & \times \mathcal{I}_\alpha(w'_1, \eta'_1, \epsilon'_1, 0, \ell_a, 0, \ell_1, 0) \prod_{i \in \{2, 3, 4\}} \left(\ell_i - w_i, \ell_i - 2w_i, \eta_i, \epsilon_i, w_i - \eta_i - \epsilon_i \right) \\
 & \times \mathcal{I}_\beta(x'_{i,12}, x'_{j,12}, x'_{k,12}, 2\eta'_1, 2\eta_2) \\
 & \times \mathcal{I}_\beta(y'_{i,12}, y'_{j,12}, y'_{k,12}, 2\epsilon'_1, 2\epsilon_2) \\
 & \times \mathcal{I}_\beta(z'_{i,12}, 0, z'_{k,12}, \ell'_1 - 2\eta'_1 - 2\epsilon'_1, \ell_2 - 2\eta_2 - 2\epsilon_2) \\
 & \times \mathcal{I}_\beta(x_{i,34}, x_{j,34}, x_{k,34}, 2\eta_3, 2\eta_4) \\
 & \times \mathcal{I}_\beta(y_{i,34}, y_{j,34}, y_{k,34}, 2\epsilon_3, 2\epsilon_4) \\
 & \times \mathcal{I}_\beta(z_{i,34}, 0, z_{k,34}, \ell_3 - 2\eta_3 - 2\epsilon_3, \ell_4 - 2\eta_4 - 2\epsilon_4) \\
 & \times \mathcal{I}_\gamma(f'_1, n_a, \ell_a, n_1, \ell_1) \left[\prod_{i \in \{2, 3, 4\}} \binom{n_i + 2\ell_i + 2}{n_i - f_i} \frac{1}{f_i!} \right] \\
 & \times \mathcal{I}_\beta(t'_{12}, 0, 0, f'_1, f_2) \mathcal{I}_\beta(t_{34}, 0, 0, f_3, f_4) \\
 & \times \mathcal{S}_A(t'_{12}, r'_{k,12}, r'_{i,12}, r'_{j,12}, t_{34}, r_{k,34}, r_{i,34}, r_{j,34}, \ell_s) \\
 & \times (-1)^{w_s - x_{i,s} - y_{i,s} - y_{j,s} + f_s} \frac{1}{2^{\ell_s + 10}} \frac{1}{\pi^4 \sqrt{\pi}} \Lambda^{(0)} 4^{-6 - \ell_s - f_s} \\
 & \times \frac{(t'_{12} + r'_{k,12} - 1)!! (r'_{i,12} - 1)!! (r'_{j,12} - 1)!! (t_{34} + r_{k,34} - 1)!! (r_{i,34} - 1)!! (r_{j,34} - 1)!!}{(t'_{12} + r'_{i,12} + r'_{j,12} + r'_{k,12} + 1)!! (t_{34} + r_{i,34} + r_{j,34} + r_{k,34} + 1)!!} \\
 & \times \frac{(5 + \ell_s + f_s)! (r_{i,s} + r_{j,s} + 2)!! (1 - r_{i,s} - r_{j,s} - r_{k,s} + \ell_s + t_s)!!}{(5 - r_{k,s} + \ell_s + t_s)!!} \\
 & \times \frac{(z_{i,s} + 2\eta_s - x_{i,s} - x_{j,s} + 2\epsilon_s - y_{i,s} - y_{j,s})!! (x_{i,s} + y_{i,s} + \ell_s - 2\epsilon_s - 2\eta_s - z_{i,s} - 1)!!}{(1 + \ell_s - x_{j,s} - y_{j,s})!!} \\
 & \times \frac{(x_{j,s} + 2\epsilon_s - y_{j,s} - 1)!! (y_{j,s} + 2\eta_s - x_{j,s} - 1)!!}{(2\epsilon_s + 2\eta_s)!!},
 \end{aligned}$$

**A: 34 Fold
Finite sum**

← **massless
particles** →

$$\begin{aligned}
 & C_{n_a \ell_a m_a, n_b \ell_b m_b, n_1 \ell_1 m_1, n_2 \ell_2 m_2, n_3 \ell_3 m_3, n_4 \ell_4 m_4} \\
 & = \prod_{i \in \{a, b\}} \frac{n_i!}{\Gamma(2\ell_i + n_i + 3)} \prod_{i \in \{a, b, 1, 2, 3, 4\}} \sqrt{2\ell_i + 1} \left[\frac{(\ell_i - |m_i|)!}{(\ell_i + |m_i|)!} \right]^{1/2} \mathcal{F}_\alpha(m_i) \\
 & \sum_{w'_i=0}^{\lfloor \frac{\ell_a - |m_a|}{2} \rfloor + \lfloor \frac{\ell_1 - |m_1|}{2} \rfloor} \sum_{p'_i=0}^{|m_a| + |m_1|} \sum_{w'_i=0}^{\lfloor \frac{\ell_b - |m_b|}{2} \rfloor + \lfloor \frac{\ell_2 - |m_2|}{2} \rfloor} \sum_{p'_i=0}^{|m_b| + |m_2|} \sum_{f'_i=0}^{n_a + n_1} \sum_{f'_i=0}^{n_b + n_2} \left[\prod_{i \in \{1, 2\}} \sum_{\eta'_i=0}^{w'_i - \eta'_i} \sum_{\epsilon'_i=0}^{f'_i + f'_2} \right] \sum_{t''_{12}=0}^{f'_3 + f'_4} \\
 & \left[\prod_{i \in \{3, 4\}} \sum_{w_i=0}^{\lfloor \frac{\ell_i - |m_i|}{2} \rfloor} \sum_{p_i=0}^{|m_i|} \sum_{\eta_i=0}^{w_i - \eta_i} \sum_{\epsilon_i=0}^{p_i} \sum_{f_i=0}^{n_i} \right] \sum_{t_{34}=0}^{f_3 + f_4} \\
 & \sum_{x''_{i,12} + x''_{j,12} + x''_{k,12} \leq 2\eta''_{12} + p'_{12}} \sum_{y''_{i,12} + y''_{j,12} + y''_{k,12} \leq 2\epsilon''_{12} + |m''_{12} - p'_{12}|} \sum_{z''_{i,12} + z''_{k,12} \leq \ell''_{12} - |m''_{12} - 2\eta''_{12} - 2\epsilon''_{12}|} \\
 & \sum_{x''_{i,12}, x''_{j,12}, x''_{k,12} = 0} \sum_{y''_{i,12}, y''_{j,12}, y''_{k,12} = 0} \sum_{z''_{i,12}, z''_{k,12} = 0} \\
 & \sum_{x_{i,34} + x_{j,34} + x_{k,34} \leq 2\eta_{34} + p_{34}} \sum_{y_{i,34} + y_{j,34} + y_{k,34} \leq 2\epsilon_{34} + |m_{34} - p_{34}|} \sum_{z_{i,34} + z_{k,34} \leq \ell_{34} - |m_{34} - 2\eta_{34} - 2\epsilon_{34}|} \\
 & \sum_{x_{i,34}, x_{j,34}, x_{k,34} = 0} \sum_{y_{i,34}, y_{j,34}, y_{k,34} = 0} \sum_{z_{i,34}, z_{k,34} = 0} \\
 & \times \mathcal{I}_\alpha(w'_1, \eta'_1, \epsilon'_1, p'_1, \ell_a, m_a, \ell_1, m_1) \mathcal{I}_\alpha(w'_2, \eta'_2, \epsilon'_2, p'_2, \ell_b, m_b, \ell_2, m_2) \\
 & \prod_{i \in \{3, 4\}} \left(\ell_i - w_i, \ell_i - 2w_i - |m_i|, p_i, |m_i| - p_i, \eta_i, \epsilon_i, w_i - \eta_i - \epsilon_i \right) |m_i|! e^{\mathcal{F}_\beta(m_i) \times i((|m_i| - p_i) \frac{\pi}{2})} \\
 & \times \mathcal{I}_\beta(x''_{i,12}, x''_{j,12}, x''_{k,12}, 2\eta'_1 + p'_1, 2\eta'_2 + p'_2) \\
 & \times \mathcal{I}_\beta(y''_{i,12}, y''_{j,12}, y''_{k,12}, 2\epsilon'_1 + |m'_1| - p'_1, 2\epsilon'_2 + |m'_2| - p'_2) \\
 & \times \mathcal{I}_\beta(z''_{i,12}, 0, z''_{k,12}, \ell'_1 - |m'_1| - 2\eta'_1 - 2\epsilon'_1, \ell'_2 - |m'_2| - 2\eta'_2 - 2\epsilon'_2) \\
 & \times \mathcal{I}_\beta(x_{i,34}, x_{j,34}, x_{k,34}, 2\eta_3 + p_3, 2\eta_4 + p_4) \\
 & \times \mathcal{I}_\beta(y_{i,34}, y_{j,34}, y_{k,34}, 2\epsilon_3 + |m_3| - p_3, 2\epsilon_4 + |m_4| - p_4) \\
 & \times \mathcal{I}_\beta(z_{i,34}, 0, z_{k,34}, \ell_3 - |m_3| - 2\eta_3 - 2\epsilon_3, \ell_4 - |m_4| - 2\eta_4 - 2\epsilon_4) \\
 & \times \mathcal{I}_\gamma(f'_1, n_a, \ell_a, n_1, \ell_1) \mathcal{I}_\gamma(f'_2, n_b, \ell_b, n_2, \ell_2) \left[\prod_{i \in \{3, 4\}} \binom{n_i + 2\ell_i + 2}{n_i - f_i} \frac{1}{f_i!} \right] \\
 & \times \mathcal{I}_\beta(t''_{12}, 0, 0, f'_1, f'_2) \mathcal{I}_\beta(t_{34}, 0, 0, f_3, f_4) \\
 & \times \mathcal{S}_C(t''_{12}, r''_{k,12}, r''_{i,12}, r''_{j,12}, t_{34}, r_{k,34}, r_{i,34}, r_{j,34}, p_s, |m|_s, \ell_s) \\
 & \times (-1)^{w_s - x_{i,s} - y_{i,s} - y_{j,s} + f_s} \frac{1}{2^{\ell_s + 9}} \frac{1}{\pi^2} \frac{1}{\Lambda^2} \sigma^{(0)} 4^{-6 - \ell_s - f_s} \\
 & \times \frac{(t''_{12} + r''_{k,12} - 1)!! (r''_{i,12} - 1)!! (r''_{j,12} - 1)!! (t_{34} + r_{k,34} - 1)!! (r_{i,34} - 1)!! (r_{j,34} - 1)!!}{(t''_{12} + r''_{i,12} + r''_{j,12} + r''_{k,12} + 1)!! (t_{34} + r_{i,34} + r_{j,34} + r_{k,34} + 1)!!} \\
 & \times \frac{(5 + \ell_s + f_s)! (r_{i,s} + r_{j,s} + 2)!! (1 - r_{i,s} - r_{j,s} - r_{k,s} + \ell_s + t_s)!!}{(5 - r_{k,s} + \ell_s + t_s)!!} \\
 & \times \frac{(z_{i,s} + 2\eta_s - x_{i,s} - x_{j,s} + 2\epsilon_s + |m|_s - y_{i,s} - y_{j,s})!! (x_{i,s} + y_{i,s} + \ell_s - 2\epsilon_s - |m|_s - 2\eta_s - z_{i,s} - 1)!!}{(1 + \ell_s - x_{j,s} - y_{j,s})!!} \\
 & \times \frac{(x_{j,s} + (2\epsilon_s + |m|_s - p_s) - y_{j,s} - 1)!! (y_{j,s} + (2\eta_s + p_s) - x_{j,s} - 1)!!}{((2\epsilon_s + |m|_s - p_s) + (2\eta_s + p_s))!!},
 \end{aligned}$$

**C: 38 Fold
Finite sum**